

ST RECEPTOR BINDING COMPOUNDS AND METHODS OF USING THE SAME*Small***FIELD OF THE INVENTION**

The present invention relates to compounds which comprise a receptor ligand moiety conjugated to an active agent. More particularly, the present invention relates to compounds which comprise a moiety that binds to the ST receptor conjugated to a therapeutic or imaging moiety.

BACKGROUND OF THE INVENTION

Colorectal cancer is the third most common neoplasm worldwide and the second most common in the United States, representing about 15% of the newly diagnosed cases of cancer in the United States. The large intestine or large bowel is the third leading site for the development of new cancer and is diagnosed in about 150,000 patients each year. Colorectal cancer is the second leading cause of cancer-related deaths and is responsible for about 12% of cancer deaths in the United States. The mortality rate of newly diagnosed large bowel cancer approaches 50% and there has been little improvement over the past 40 years. Most of this mortality reflects local, regional and distant metastases. About thirty percent of patients with colorectal cancer have unresectable disease at presentation and about 40% develop metastases during the course of their disease. Distant metastatic disease is seen in liver (about 12%), lung (about 3%), bone (about 0.9%), brain (about 0.7%), nodes (about 4%), and peritoneum (about 2%) at the time of initial diagnosis. In 1987, the large bowel cancers found

09724983-112800

regionally or at distant sites at the time of diagnosis were about 26% and about 18%, respectively.

5 Surgery is the mainstay of treatment for colorectal cancer but recurrence is frequent. Colorectal cancer has been achieved using a combination of 5-fluorouracil and levamisole. Surgery has had the largest impact on survival and, in some patients with limited disease, achieves a cure. However, surgery removes bulk tumor, leaving behind microscopic residual disease which ultimately results in recrudescence. Overall recurrence rates for colonic tumors are about 33% and for rectal cancer about 42%. Of these recurrences, about 9% are local, about 13% are systemic metastatic disease, and the remaining 88% are a combination of local and systemic disease. 10 Fifty percent of patients with recurrent colorectal cancer have hepatic metastases.

Early detection of primary, metastatic, and recurrent disease can significantly impact the prognosis of individuals suffering from colorectal cancer. Large bowel cancer diagnosed at an early stage has a significantly better outcome than that diagnosed at more advanced stages. The 5 year relative survival rates for patients with regional or distant metastases are 48% and 5%, compared with 90% and 77% for disease which is *in situ* or local, respectively, at the time of diagnosis. 20 Similarly, diagnosis of metastatic or recurrent disease earlier potentially carries with it a better prognosis.

Although current radiotherapeutic agents, chemotherapeutic agents and biological toxins are potent cytotoxins, they do not discriminate between normal and malignant cells, producing adverse effects and dose-limiting toxicities. Over the past decade, a novel approach has been employed to more specifically target agents to tumor cells, involving the conjugation of an active agent to molecules which binds preferentially to antigens that exist predominantly on tumor cells. These conjugates can be administered systemically and specifically bind to the targeted tumor cells. 35 Theoretically, targeting permits uptake by cells of cytotoxic

008277 8842260

agents at concentrations which do not produce serious toxicities in normal tissues. Also, selective binding to targeted tumor cells facilitates detection of occult tumor and is therefore useful in designing imaging agents. Molecular
5 targeting predominantly has employed monoclonal antibodies generated to antigens selectively expressed on tumor cells.

Immunoscintigraphy using monoclonal antibodies directed at tumor-specific markers has been employed to diagnose colorectal cancer. Monoclonal antibodies against
10 carcinoembryonic antigen (CEA) labeled with ⁹⁹Technetium identified 94% of patients with recurrent tumors. Similarly, ¹¹¹Indium-labeled anti-CEA monoclonal antibodies successfully diagnosed 85% of patients with recurrent colorectal carcinoma who were not diagnosed by conventional techniques. ¹²⁵Iodine-
15 labeled antibodies have been effective in localizing more than 80% of the pathologically-confirmed recurrences by intraoperative gamma probe scanning.

Monoclonal antibodies have also been employed to target specific therapeutic agents in colorectal cancer.
20 Preclinical studies demonstrated that anti-CEA antibodies labeled with ⁹⁰Yttrium inhibited human colon carcinoma xenografts in nude mice. Antibodies generated to colorectal cancer cells and coupled to mitomycin C or neocarzinostatin demonstrated an anti-tumor effect on human colon cancer
25 xenografts in nude mice and 3 patients with colon cancer. Similar results in animals were obtained with monoclonal antibodies conjugated to ricin toxin A chain.

Due to the sensitivity, specificity, and adverse-effect profile of monoclonal antibodies, the results obtained
30 using monoclonal antibody-based therapeutics have shown them to be less than ideal targeting tools. Although monoclonal antibodies have been generated to antigens selectively expressed on tumors, no truly cancer-specific antibody has been identified. Most antigens expressed on neoplastic cells appear
35 to be quantitatively increased in these compared to normal cells but the antigens are nonetheless often present in normal cells. Thus, antibodies to such determinants can react with

09724983 "112800

non-neoplastic tissues, resulting in significant toxicities. Also, antibodies are relatively large molecules and consequently, often evoke an immune response in patients. These immune responses can result in significant toxicities in
5 patients upon re-exposure to the targeting agents and can prevent targeting by the monoclonal due to immune complex formation with degradation and excretion. Finally, binding of antibodies to tumor cells may be low and targeted agents may be delivered to cells in quantities insufficient to achieve
10 detection or cytotoxicity.

There remains a need for compositions which can specifically target metastasized colorectal cancer cells. There is a need for imaging agents which can specifically bind to metastasized colorectal cancer cells. There is a need for
15 improved methods of imaging metastasized colorectal cancer cells. There is a need for therapeutic agents which can specifically bind to metastasized colorectal cancer cells. There is a need for improved methods of treating individuals who are suspected of suffering from colorectal cancer cells,
20 especially individuals who are suspected of suffering from metastasis of colorectal cancer cells.

SUMMARY OF THE INVENTION

The present invention relates to conjugated compounds which comprises an ST receptor binding moiety and a radiostable
25 active moiety.

The present invention relates to a pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent, and a conjugated compound which comprises an ST receptor binding moiety and a radiostable active moiety.

30 The present invention relates to a method of treating an individual suspected of suffering from metastasized colorectal cancer comprising the steps of administering to said individual a pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent, and a
35 therapeutically effective amount of a conjugated compound which

00827T " E864260

comprises an ST receptor binding moiety and a radiostable active moiety.

The present invention relates to a pharmaceutical composition comprising a pharmaceutically acceptable carrier or
5 diluent, and conjugated compound that comprises an ST receptor binding moiety and a radioactive active moiety wherein the conjugated compound is present in an amount effective for therapeutic or diagnostic use in humans suffering from colorectal cancer.

10 The present invention relates to a method of radioimaging metastasized colorectal cancer cells comprising the steps of first administering to an individual suspected of having metastasized colorectal cancer cells, a pharmaceutical composition that comprises a pharmaceutically acceptable
15 carrier or diluent, and conjugated compound that comprises an ST receptor binding moiety and a radioactive active moiety wherein the conjugated compound is present in an amount effective for diagnostic use in humans suffering from colorectal cancer and then detecting the localization and
20 accumulation of radioactivity in the individual's body.

The present invention relates to a method of treating an individual suspected of suffering from metastasized colorectal cancer comprising the steps of administering to said individual a pharmaceutical composition comprising a
25 pharmaceutically acceptable carrier or diluent, and a therapeutically effective amount of a conjugated compound which comprises an ST receptor binding moiety and a radioactive active moiety.

DESCRIPTION OF PREFERRED EMBODIMENTS OF THE INVENTION

30 As used herein, the terms "ST" and "native ST" are used interchangeably and are meant to refer to heat-stable toxin (ST) which is a peptide produced by *E. coli*, as well as other organisms. STs are naturally occurring peptides which 1) are naturally produced by organisms, 2) which bind to the ST
35 receptor and 3) which activate the signal cascade that mediates ST-induced diarrhea.

008211 112800

As used herein, the term "ST receptor" is meant to refer to the receptors found on colorectal cells, including local and metastasized colorectal cancer cells, which bind to ST. In normal individuals, ST receptors are found exclusively
5 in cells of intestine, in particular in cells in the duodenum, small intestine (jejunum and ileum), the large intestine, colon (cecum, ascending colon, transverse colon, descending colon and sigmoid colon) and rectum.

As used herein, the term "ST receptor ligand" is meant
10 to refer to compounds which specifically bind to the ST receptor. ST is an ST receptor ligand. An ST receptor ligand may be a peptide or a non-peptide.

As used herein, the term "ST receptor binding peptide" is meant to refer to ST receptor ligands that are peptides.

15 As used herein, the term "ST peptides" is meant to refer to ST receptor binding peptides selected from the group consisting of SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof.

As used herein, the term "fragment" is meant to refer
20 to peptide a) which has an amino acid sequence identical to a portion of an ST receptor binding peptide and b) which is capable of binding to the ST receptor.

As used herein, the term "derivative" is meant to refer to a peptide a) which has an amino acid sequence
25 substantially identical to at least a portion of an ST receptor binding peptide and b) which is capable of binding to the ST receptor.

As used herein, the term "substantially identical" is meant to refer to an amino acid sequence that is the same as
30 the amino acid sequence of an ST peptide except some of the residues are deleted or substituted with conservative amino acids or additional amino acids are inserted.

As used herein, the term "active agent" is meant to refer to compounds that are therapeutic agents or imaging
35 agents.

009724983 112800

As used herein, the term "radiostable" is meant to refer to compounds which do not undergo radioactive decay; i.e. compounds which are not radioactive.

As used herein, the term "therapeutic agent" is meant
5 to refer to chemotherapeutics, toxins, radiotherapeutics, targeting agents or radiosensitizing agents.

As used herein, the term "chemotherapeutic" is meant to refer to compounds that, when contacted with and/or incorporated into a cell, produce an effect on the cell
10 including causing the death of the cell, inhibiting cell division or inducing differentiation.

As used herein, the term "toxin" is meant to refer to compounds that, when contacted with and/or incorporated into a cell, produce the death of the cell.

As used herein, the term "radiotherapeutic" is meant
15 to refer to radionuclides which when contacted with and/or incorporated into a cell, produce the death of the cell.

As used herein, the term "targeting agent" is meant to refer compounds which can be bound by and or react with
20 other compounds. Targeting agents may be used to deliver chemotherapeutics, toxins, enzymes, radiotherapeutics, antibodies or imaging agents to cells that have targeting agents associated with them and/or to convert or otherwise transform or enhance coadministered active agents. A targeting
25 agent may include a moiety that constitutes a first agent that is localized to the cell which when contacted with a second agent either is converted to a third agent which has a desired activity or causes the conversion of the second agent into an agent with a desired activity. The result is the localized
30 agent facilitates exposure of an agent with a desired activity to the metastasized cell.

As used herein, the term "radiosensitizing agent" is meant to refer to agents which increase the susceptibility of cells to the damaging effects of ionizing radiation. A
35 radiosensitizing agent permits lower doses of radiation to be administered and still provide a therapeutically effective dose.

008211 88542250

As used herein, the term "imaging agent" is meant to refer to compounds which can be detected.

As used herein, the term "ST receptor binding moiety" is meant to refer to the portion of a conjugated compound that
5 constitutes an ST receptor ligand.

As used herein, the term "active moiety" is meant to refer to the portion of a conjugated compound that constitutes an active agent.

As used herein, the terms "conjugated compound" and
10 "conjugated composition" are used interchangeably and meant to refer to a compound which comprises an ST receptor binding moiety and an active moiety and which is capable of binding to the ST receptor. Conjugated compounds according to the present invention comprise a portion which constitutes an ST receptor
15 ligand and a portion which constitutes an active agent. Thus, conjugated compounds according to the present invention are capable of specifically binding to the ST receptor and include a portion which is a therapeutic agent or imaging agent. Conjugated compositions may comprise crosslinkers and/or
20 molecules that serve as spacers between the moieties.

As used herein, the terms "crosslinker", "crosslinking agent", "conjugating agent", "coupling agent", "condensation reagent" and "bifunctional crosslinker" are used interchangeably and are meant to refer to molecular groups
25 which are used to attach the ST receptor ligand and the active agent to thus form the conjugated compound.

As used herein, the term "colorectal cancer" is meant to include the well-accepted medical definition that defines colorectal cancer as a medical condition characterized by
30 cancer of cells of the intestinal tract below the small intestine (i.e. the large intestine (colon), including the cecum, ascending colon, transverse colon, descending colon, and sigmoid colon, and rectum). Additionally, as used herein, the term "colorectal cancer" is meant to further include medical
35 conditions which are characterized by cancer of cells of the duodenum and small intestine (jejunum and ileum). The definition of colorectal cancer used herein is more expansive

003211 6864260

than the common medical definition but is provided as such since the cells of the duodenum and small intestine also contain ST receptors and are therefore amenable to the methods of the present invention using the compounds of the present invention.

As used herein, the term "metastasis" is meant to refer to the process in which cancer cells originating in one organ or part of the body relocate to another part of the body and continue to replicate. Metastasized cells subsequently form tumors which may further metastasize. Metastasis thus refers to the spread of cancer from the part of the body where it originally occurs to other parts of the body. The present invention relates to methods of delivering active agents to metastasized colorectal cancer cells.

As used herein, the term "metastasized colorectal cancer cells" is meant to refer to colorectal cancer cells which have metastasized; colorectal cancer cells localized in a part of the body other than the duodenum, small intestine (jejunum and ileum), large intestine (colon), including the cecum, ascending colon, transverse colon, descending colon, and sigmoid colon, and rectum.

ST, which is produced by *E. coli*, as well as other organisms, is responsible for endemic diarrhea in developing countries and travelers diarrhea. ST induces intestinal secretion by binding to specific receptors, ST receptors, in the apical brush border membranes of the mucosal cells lining the intestinal tract. Binding of ST to ST receptors is non-covalent and occurs in a concentration-dependent and saturable fashion. Once bound, ST-ST receptor complexes appear to be internalized by intestinal cells, i.e. transported from the surface into the interior of the cell. Binding of ST to ST receptors triggers a cascade of biochemical reactions in the apical membrane of these cells resulting in the production of a signal which induces intestinal cells to secrete fluids and electrolytes, resulting in diarrhea.

ST receptors are unique in that they are only localized in the apical brush border membranes of the cells

09724983-112800

lining the intestinal tract. Indeed, they are not found in any other cell type in placental mammals. In addition, ST receptors are almost exclusively localized to the apical membranes, with little being found in the basolateral membranes on the sides of intestinal cells.

Mucosal cells lining the intestine are joined together by tight junctions which form a barrier against the passage of intestinal contents into the blood stream and components of the blood stream into the intestinal lumen. Therefore, the apical location of ST receptors isolates these receptors from the circulatory system so that they may be considered to exist separate from the rest of the body; essentially the "outside" of the body. Therefore, the rest of the body is considered "outside" the intestinal tract. Compositions administered "outside" the intestinal tract are maintained apart and segregated from the only cells which normally express ST receptors.

In individuals suffering from colorectal cancer, the cancer cells are often derived from cells that produce and display the ST receptor and these cancer cells continue to produce and display the ST receptor on their cell surfaces. Indeed, T84 cells, which are human colonic adenocarcinoma cells isolated from lung metastases, express ST receptors on their cell surface. Similarly, HT29glu-cells, which are human colonic adenocarcinoma cells, express receptors for ST. Thus, in individuals suffering from colorectal cancer, some metastasized intestinal cancer cells express ST receptors.

An effort was undertaken to determine the proportion of colorectal tumors which have the ST receptor. Sixteen colorectal cancer tumors, including ten local colorectal tumors and six metastasized tumors (3 liver, 1 lung, 1 lymphnode, 1 peritoneum), were tested and each possessed ST receptors. In each case, the affinity and density of receptors was amenable for targeting. That is, the cells possessed at least 10^4 - 10^6 receptors per cell and demonstrated an affinity of 10^{-7} or better (that is preferably between 10^{-8} to 10^{-9} or less; the lower number indicating a tighter bond, thus a higher

05724983 112800

affinity). Normal liver, lymphnode, peritoneum and lung cells were found not to possess ST receptors.

When such cancer cells metastasize, the metastasized cancer cells continue to produce and display the ST receptor.

5 The expression of ST receptors on the surfaces of metastatic tumors provides a target for selective binding of conjugated compositions. ST receptors permit the absolutely specific targeting of therapeutic and diagnostic agents that are conjugated to ST receptor ligands to metastatic colorectal

10 cancer cells.

The conjugated compositions of the present invention are useful for targeting cells that line the inner intestine wall including those cancer cells derived from such cells, particularly metastasized cancer cells derived from such cells.

15 The conjugated compositions of the present invention which are administered outside the intestinal tract such as those administered in the circulatory system will remain segregated from the cells that line the intestinal tract and will bind only to cells outside the intestinal tract which are derived

20 from the intestinal tract such as metastasized colorectal cells. The conjugated compositions will not bind to non-colorectal derived cells. Thus, the active moieties of conjugated compositions administered outside the intestinal tract are delivered to cells which are derived from the

25 intestinal tract such as metastasized colorectal cells but will not be delivered to any other cells.

Therapeutic and diagnostic pharmaceutical compositions of the present invention include conjugated compounds specifically targeted to metastatic disease. These conjugated

30 compounds include ST receptor binding moieties which do not bind to cells of normal tissue in the body except cells of the intestinal tract since the cells of other tissues do not possess ST receptors. Unlike normal colorectal cells and localized colorectal cancer cells, metastasized colorectal

35 cancer cells are accessible to substances administered outside the intestinal tract, for example administered in the circulatory system. The only ST receptors in normal tissue

09724983 112800

exist in the apical membranes of intestinal mucosa cells and these receptors are effectively isolated from the targeted cancer chemotherapeutics and imaging agents administered outside the intestinal tract by the intestinal mucosa barrier.

5 Thus, metastasized colorectal cells may be targeted by conjugated compounds of the present invention by introducing such compounds outside the intestinal tract such as for example by administering pharmaceutical compositions that comprise conjugated compounds into the circulatory system.

10 One having ordinary skill in the art can readily
identify individuals suspected of suffering from colorectal
cancer and metastasized colorectal cells. In those individuals
diagnosed with colorectal cancer, it is standard therapy to
suspect metastasis and aggressively attempt to eradicate
15 metastasized cells. The present invention provides
pharmaceutical compositions and methods for imaging and thereby
will more definitively diagnose metastasis. Further, the
present invention provides pharmaceutical compositions
comprising therapeutic agents and methods for specifically
20 targeting and eliminating metastasized colorectal cancer cells.
Further, the present invention provides pharmaceutical
compositions that comprise therapeutics and methods for
specifically eliminating colorectal cancer cells.

The pharmaceutical compositions which comprise conjugated compositions of the present invention may be used to diagnose or treat individuals suffering from localized colorectal tumors, that is primary or non-metastatic colorectal tumors if these have penetrated the basement membrane underlying the mucosa into the submucosa where there is abundant blood supply to which they have access. Penetration into the submucosa circumvents the mucosal barrier resulting in the ability of conjugated compositions introduced into the circulatory system to interact with these tumors.

The present invention relies upon the use of an ST
35 receptor binding moiety in a conjugated composition. The ST
receptor binding moiety is essentially a portion of the
conjugated composition which acts as a ligand to the ST

receptor and thus specifically binds to these receptors. The conjugated composition also includes an active moiety which is associated with the ST receptor binding moiety; the active moiety being an active agent which is either useful to image, target, neutralize or kill the cell.

According to the present invention, the ST receptor binding moiety is the ST receptor ligand portion of a conjugated composition. In some embodiments, the ST receptor ligand may be native ST.

Native ST has been isolated from a variety of organisms including *E. coli*, *Yersinia*, *Enterobacter*, and others. In nature, the toxins are generally encoded on a plasmid which can "jump" between different species. Several different toxins have been reported to occur in different species. These toxins all possess significant sequence homology, they all bind to ST receptors and they all activate guanylate cyclase, producing diarrhea.

ST has been both cloned and synthesized by chemical techniques. The cloned or synthetic molecules exhibit binding characteristics which are similar to native ST. Native ST isolated from *E. coli* is 18 or 19 amino acids in length. The smallest "fragment" of ST which retains activity is the 13 amino acid core peptide extending toward the carboxy terminal from cysteine 6 to cysteine 18 (of the 19 amino acid form). Analogues of ST have been generated by cloning and by chemical techniques. Small peptide fragments of the native ST structure which include the structural determinant that confers binding activity may be constructed. Once a structure is identified which binds to ST receptors, non-peptide analogues mimicking that structure in space are designed.

SEQ ID NO:1 discloses a nucleotide sequence which encodes 19 amino acid ST, designated ST Ia, reported by So and McCarthy (1980) *Proc. Natl. Acad. Sci. USA* 77:4011, which is incorporated herein by reference.

The amino acid sequence of ST Ia is disclosed in SEQ ID NO:2.

09724983 112800

SEQ ID NO:3 discloses the amino acid sequence of an 18 amino acid peptide which exhibits ST activity, designated ST I*, reported by Chan and Giannella (1981) *J. Biol. Chem.* 256:7744, which is incorporated herein by reference.

5 SEQ ID NO:4 discloses a nucleotide sequence which encodes 19 amino acid ST, designated ST Ib, reported by Mosely et al. (1983) *Infect. Immun.* 39:1167, which is incorporated herein by reference.

The amino acid sequence of ST Ib is disclosed in SEQ
10 ID NO:5.

A 15 amino acid peptide called guanylin which has about 50% sequence homology to ST has been identified in mammalian intestine (Currie, M.G. et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:947-951, which is incorporated herein by
15 reference). Guanylin binds to ST receptors and activates guanylate cyclase at a level of about 10- to 100-fold less than native ST. Guanylin may not exist as a 15 amino acid peptide in the intestine but rather as part of a larger protein in that organ. The amino acid sequence of guanylin from rodent is
20 disclosed as SEQ ID NO:6.

SEQ ID NO:7 is an 18 amino acid fragment of SEQ ID NO:2. SEQ ID NO:8 is a 17 amino acid fragment of SEQ ID NO:2. SEQ ID NO:9 is a 16 amino acid fragment of SEQ ID NO:2. SEQ ID NO:10 is a 15 amino acid fragment of SEQ ID NO:2. SEQ ID NO:11
25 is a 14 amino acid fragment of SEQ ID NO:2. SEQ ID NO:12 is a 13 amino acid fragment of SEQ ID NO:2. SEQ ID NO:13 is an 18 amino acid fragment of SEQ ID NO:2. SEQ ID NO:14 is a 17 amino acid fragment of SEQ ID NO:2. SEQ ID NO:15 is a 16 amino acid fragment of SEQ ID NO:2. SEQ ID NO:16 is a 15 amino acid
30 fragment of SEQ ID NO:2. SEQ ID NO:17 is a 14 amino acid fragment of SEQ ID NO:2.

SEQ ID NO:18 is a 17 amino acid fragment of SEQ ID NO:3. SEQ ID NO:19 is a 16 amino acid fragment of SEQ ID NO:3. SEQ ID NO:20 is a 15 amino acid fragment of SEQ ID NO:3. SEQ
35 ID NO:21 is a 14 amino acid fragment of SEQ ID NO:3. SEQ ID NO:22 is a 13 amino acid fragment of SEQ ID NO:3. SEQ ID NO:23 is a 17 amino acid fragment of SEQ ID NO:3. SEQ ID NO:24 is a

008211 112800

16 amino acid fragment of SEQ ID NO:3. SEQ ID NO:25 is a 15 amino acid fragment of SEQ ID NO:3. SEQ ID NO:26 is a 14 amino acid fragment of SEQ ID NO:3.

SEQ ID NO:27 is an 18 amino acid fragment of SEQ ID NO:5. SEQ ID NO:28 is a 17 amino acid fragment of SEQ ID NO:5. SEQ ID NO:29 is a 16 amino acid fragment of SEQ ID NO:5. SEQ ID NO:30 is a 15 amino acid fragment of SEQ ID NO:5. SEQ ID NO:31 is a 14 amino acid fragment of SEQ ID NO:5. SEQ ID NO:32 is a 13 amino acid fragment of SEQ ID NO:5. SEQ ID NO:33 is an 18 amino acid fragment of SEQ ID NO:5. SEQ ID NO:34 is a 17 amino acid fragment of SEQ ID NO:5. SEQ ID NO:35 is a 16 amino acid fragment of SEQ ID NO:5. SEQ ID NO:36 is a 15 amino acid fragment of SEQ ID NO:5. SEQ ID NO:37 is a 14 amino acid fragment of SEQ ID NO:5.

SEQ ID NO:27, SEQ ID NO:31, SEQ ID NO:36 AND SEQ ID NO:37 are disclosed in Yoshimura, S., et al. (1985) *FEBS Lett.* 181:138, which is incorporated herein by reference.

SEQ ID NO:38, SEQ ID NO:39 and SEQ ID NO:40, which are derivatives of SEQ ID NO:3, are disclosed in Waldman, S.A. and O'Hanley, P. (1989) *Infect. Immun.* 57:2420, which is incorporated herein by reference.

SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44 and SEQ ID NO:45, which are a derivatives of SEQ ID NO:3, are disclosed in Yoshimura, S., et al. (1985) *FEBS Lett.* 181:138, which is incorporated herein by reference.

SEQ ID NO:46 is a 25 amino acid peptide derived from *Y. enterocolitica* which binds to the ST receptor.

SEQ ID NO:47 is a 16 amino acid peptide derived from *V. cholerae* which binds to the ST receptor. SEQ ID NO:47 is reported in Shimonishi, Y., et al. *FEBS Lett.* 215:165, which is incorporated herein by reference.

SEQ ID NO:48 is an 18 amino acid peptide derived from *Y. enterocolitica* which binds to the ST receptor. SEQ ID NO:48 is reported in Okamoto, K., et al. *Infec. Immun.* 55:2121, which is incorporated herein by reference.

SEQ ID NO:49, is a derivative of SEQ ID NO:5.

05724983 112800

SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53 are derivatives.

SEQ ID NO:54 is the amino acid sequence of guanylin from human.

5 In some preferred embodiments, conjugated compounds comprise ST receptor binding moieties that comprise amino acid sequences selected from the group consisting of SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof.

10 Those having ordinary skill in the art can readily design and produce derivatives having substantially identical amino acid sequences of ST peptides with deletions and/or insertions and/or conservative substitutions of amino acids. For example, following what are referred to as Dayhof's rules
15 for amino acid substitution (Dayhof, M.D. (1978) *Nat. Biomed. Res. Found.*, Washington, D.C. Vol. 5, supp. 3), amino acid residues in a peptide sequence may be substituted with comparable amino acid residues. Such substitutions are well-known and are based the upon charge and structural
20 characteristics of each amino acid. Derivatives include fragments of ST receptor binding peptides with deletions and/or insertions and/or conservative substitutions.

In some embodiments, ST receptor binding peptides comprise D amino acids. As used herein, the term "D amino acid
25 peptides" is meant to refer to ST receptor binding peptides, fragments or derivatives which comprise at least one and preferably a plurality of D amino acids which are capable of binding to the ST receptor. The use of D amino acid peptides is desirable as they are less vulnerable to degradation and
30 therefore have a longer half-life. D amino acid peptides comprising mostly all or consisting of D amino acids may comprise amino acid sequences in the reverse order of ST receptor binding peptides which are made up of L amino acids.

In some embodiments, ST receptor binding peptides,
35 including D amino acid peptides, are conformationally restricted to present and maintain the proper structural conformation for binding to the ST receptor. The compositions

09724983.112800

may comprise additional amino acid residues required to achieve proper three dimensional conformation including residues which facilitate circularization or desired folding.

It is preferred that the ST receptor ligand used as the ST receptor binding moiety be as small as possible. Thus it is preferred that the ST receptor ligand be a non-peptide small molecule or small peptide, preferably less than 25 amino acids, more preferably less than 20 amino acids. In some embodiments, the ST receptor ligand which constitute the ST receptor binding moiety of a conjugated composition is less than 15 amino acids. ST receptor binding peptide comprising less than 10 amino acids and ST receptor binding peptide less than 5 amino acids may be used as ST binding moieties according to the present invention. It is within the scope of the present invention to include larger molecules which serve as ST receptor binding moieties including, but not limited to molecules such as antibodies, FAbs and F(Ab)₂s which specifically bind to ST receptor.

An assay may be used to test both peptide and non-peptide compositions to determine whether or not they are ST receptor ligands or, to test conjugated compositions to determine if they possess ST receptor binding activity. Such compositions that specifically bind to ST receptors can be identified by a competitive binding assay. The competitive binding assay is a standard technique in pharmacology which can be readily performed by those having ordinary skill in the art using readily available starting materials. Competitive binding assays have been shown to be effective for identifying compositions that specifically bind to ST receptors. Briefly, the assay consists of incubating a preparation of ST receptors (e.g. intestinal membranes from rat intestine, human intestine, T84 cells) with a constant concentration (1×10^{-10} M to 5×10^{-10} M) of ¹²⁵I-ST (any ST receptor ligand such as native STs SEQ ID NO:2, SEQ ID NO:3 or SEQ ID NO:5 may be used) and a known concentration of a test compound. As a control, a duplicate preparation of ST receptors are incubated with a duplicate concentration of ¹²⁵I-ST in the absence of test compound.

09724983-112800

Assays are incubated to equilibrium (2 hours) and the amount of ^{125}I -ST bound to receptors is quantified by standard techniques. The ability of the test compound to bind to receptors is measured as its ability to prevent (compete with) the ^{125}I -ST from binding. Thus, in assays containing the test compound which bind to the receptor, there will be less radioactivity associated with the receptors. This assay, which is appropriate for determining the ability of any molecule to bind to ST receptors, is a standard competitive binding assay which can be readily employed by those having ordinary skill in the art using readily available starting materials.

ST may be isolated from natural sources using standard techniques. Additionally, ST receptor binding peptides and conjugated compositions or portions thereof which are peptides may be prepared routinely by any of the following known techniques.

ST receptor binding peptides and conjugated compositions or portions thereof which are peptides may be prepared using the solid-phase synthetic technique initially described by Merrifield, in *J. Am. Chem. Soc.*, 15:2149-2154 (1963). Other peptide synthesis techniques may be found, for example, in M. Bodanszky et al., (1976) *Peptide Synthesis*, John Wiley & Sons, 2d Ed.; Kent and Clark-Lewis in *Synthetic Peptides in Biology and Medicine*, p. 295-358, eds. Alitalo, K., et al. Science Publishers, (Amsterdam, 1985); as well as other reference works known to those skilled in the art. A summary of peptide synthesis techniques may be found in J. Stuart and J.D. Young, *Solid Phase Peptide Synthesis*, Pierce Chemical Company, Rockford, IL (1984), which is incorporated herein by reference. The synthesis of peptides by solution methods may also be used, as described in *The Proteins*, Vol. II, 3d Ed., p. 105-237, Neurath, H. et al., Eds., Academic Press, New York, NY (1976). Appropriate protective groups for use in such syntheses will be found in the above texts, as well as in J.F.W. McOmie, *Protective Groups in Organic Chemistry*, Plenum Press, New York, NY (1973), which is incorporated herein by reference. In general, these synthetic methods involve the

008211 112800

sequential addition of one or more amino acid residues or suitable protected amino acid residues to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid residue is protected by a suitable, 5 selectively removable protecting group. A different, selectively removable protecting group is utilized for amino acids containing a reactive side group, such as lysine.

Using a solid phase synthesis as an example, the protected or derivatized amino acid is attached to an inert 10 solid support through its unprotected carboxyl or amino group. The protecting group of the amino or carboxyl group is then selectively removed and the next amino acid in the sequence having the complementary (amino or carboxyl) group suitably protected is admixed and reacted with the residue already 15 attached to the solid support. The protecting group of the amino or carboxyl group is then removed from this newly added amino acid residue, and the next amino acid (suitably protected) is then added, and so forth. After all the desired amino acids have been linked in the proper sequence, any 20 remaining terminal and side group protecting groups (and solid support) are removed sequentially or concurrently, to provide the final peptide. The peptide of the invention are preferably devoid of benzylated or methylbenzylated amino acids. Such protecting group moieties may be used in the course of 25 synthesis, but they are removed before the peptides are used. Additional reactions may be necessary, as described elsewhere, to form intramolecular linkages to restrain conformation.

ST receptor binding peptides and conjugated compositions or portions thereof which are peptides may also be 30 prepared by recombinant DNA techniques. Provision of a suitable DNA sequence encoding the desired peptide permits the production of the peptide using recombinant techniques now known in the art. The coding sequence can be obtained from natural sources or synthesized or otherwise constructed using 35 widely available starting materials by routine methods. When the coding DNA is prepared synthetically, advantage can be

002493 112300

taken of known codon preferences of the intended host where the DNA is to be expressed.

To produce an ST receptor binding peptide which occurs in nature, one having ordinary skill in the art can, using
5 well-known techniques, obtain a DNA molecule encoding the ST receptor binding peptides from the genome of the organism that produces the ST receptor binding peptide and insert that DNA molecule into a commercially available expression vector for use in well-known expression systems.

10 Likewise, one having ordinary skill in the art can, using well known techniques, combine a DNA molecule that encodes an ST receptor binding peptide, such as SEQ ID NO:1 and SEQ ID NO:4, which can be obtained from the genome of the organism that produces the ST, with DNA that encodes a toxin,
15 another active agent that is a peptide or additionally, any other amino acid sequences desired to be adjacent to the ST receptor binding peptide amino acid sequence in a single peptide and insert that DNA molecule into a commercially available expression vector for use in well-known expression
20 systems.

For example, the commercially available plasmid pSE420 (Invitrogen, San Diego, CA) may be used for recombinant production in *E. coli*. The commercially available plasmid pYES2 (Invitrogen, San Diego, CA) may be used for production in
25 *S. cerevisiae* strains of yeast. The commercially available MaxBac™ (Invitrogen, San Diego, CA) complete baculovirus expression system may be used for production in insect cells. The commercially available plasmid pcDNA I (Invitrogen, San Diego, CA) may be used for production in mammalian cells such
30 as Chinese Hamster Ovary cells.

One having ordinary skill in the art may use these or other commercially available expression vectors and systems or produce vectors using well-known methods and readily available starting materials. Expression systems containing the
35 requisite control sequences, such as promoters and polyadenylation signals, and preferably enhancers, are readily available and known in the art for a variety of hosts. See

002493 11200

e.g., Sambrook et al., *Molecular Cloning a Laboratory Manual*, Second Ed. Cold Spring Harbor Press (1989). Thus, the desired proteins can be prepared in both prokaryotic and eukaryotic systems, resulting in a spectrum of processed forms of the protein.

The most commonly used prokaryotic system remains *E. coli*, although other systems such as *B. subtilis* and *Pseudomonas* are also useful. Suitable control sequences for prokaryotic systems include both constitutive and inducible promoters including the *lac* promoter, the *trp* promoter, hybrid promoters such as *tac* promoter, the *lambda* phage P1 promoter. In general, foreign proteins may be produced in these hosts either as fusion or mature proteins. When the desired sequences are produced as mature proteins, the sequence produced may be preceded by a methionine which is not necessarily efficiently removed. Accordingly, the peptides and proteins claimed herein may be preceded by an N-terminal Met when produced in bacteria. Moreover, constructs may be made wherein the coding sequence for the peptide is preceded by an operable signal peptide which results in the secretion of the protein. When produced in prokaryotic hosts in this matter, the signal sequence is removed upon secretion.

A wide variety of eukaryotic hosts are also now available for production of recombinant foreign proteins. As in bacteria, eukaryotic hosts may be transformed with expression systems which produce the desired protein directly, but more commonly signal sequences are provided to effect the secretion of the protein. Eukaryotic systems have the additional advantage that they are able to process introns which may occur in the genomic sequences encoding proteins of higher organisms. Eukaryotic systems also provide a variety of processing mechanisms which result in, for example, glycosylation, carboxy-terminal amidation, oxidation or derivatization of certain amino acid residues, conformational control, and so forth.

Commonly used eukaryotic systems include, but are not limited to, yeast, fungal cells, insect cells, mammalian cells,

00227T E864260

avian cells, and cells of higher plants. Suitable promoters are available which are compatible and operable for use in each of these host types as well as are termination sequences and enhancers, as e.g. the baculovirus polyhedron promoter. As
5 above, promoters can be either constitutive or inducible. For example, in mammalian systems, the mouse metallothioneine promoter can be induced by the addition of heavy metal ions.

The particulars for the construction of expression systems suitable for desired hosts are known to those in the
10 art. For recombinant production of the protein, the DNA encoding it is suitably ligated into the expression vector of choice and then used to transform the compatible host which is then cultured and maintained under conditions wherein expression of the foreign gene takes place. The protein of the
15 present invention thus produced is recovered from the culture, either by lysing the cells or from the culture medium as appropriate and known to those in the art.

One having ordinary skill in the art can, using well-known techniques, isolate the protein that is produced.

20 According to the present invention, the active moiety may be a therapeutic agent or an imaging agent. One having ordinary skill in the art can readily recognize the advantages of being able to specifically target metastasized colorectal cells with an ST receptor ligand and conjugate such a ligand
25 with many different active agents.

Chemotherapeutics useful as active moieties which when conjugated to an ST receptor binding moiety are specifically delivered to metastasized colorectal cells are typically, small chemical entities produced by chemical synthesis.
30 Chemotherapeutics include cytotoxic and cytostatic drugs. Chemotherapeutics may include those which have other effects on cells such as reversal of the transformed state to a differentiated state or those which inhibit cell replication. Examples of chemotherapeutics include common cytotoxic or
35 cytostatic drugs such as for example: methotrexate (amethopterin), doxorubicin (adrimycin), daunorubicin, cytosinarabioside, etoposide, 5-4 fluorouracil, melphalan,

09724983 112600

Ins
C1

Ins
C₁

chlorambucil, and other nitrogen mustards (e.g. cyclophosphamide), cis-platinum, vindesine (and other vinca alkaloids), mitomycin and bleomycin. Other chemotherapeutics include: purothionin (barley flour oligopeptide), macromomycin.

5 1,4-benzoquinone derivatives and trenimon.

Toxins are useful as active moieties. When a toxin is conjugated to an ST receptor binding moiety, the conjugated composition is specifically delivered to a metastasized colorectal cell by way of the ST receptor binding moiety and

10 the toxin moiety kills the cell. Toxins are generally complex toxic products of various organisms including bacteria, plants, etc. Examples of toxins include but are not limited to: ricin, ricin A chain (ricin toxin), *Pseudomonas* exotoxin (PE), diphtheria toxin (DT), *Clostridium perfringens* phospholipase C

15 (PLC), bovine pancreatic ribonuclease (BPR), pokeweed antiviral protein (PAP), abrin, abrin A chain (abin toxin), cobra venom factor (CVF), gelonin (GEL), saporin (SAP), modeccin, viscumin and volkensin. As discussed above, when protein toxins are employed with ST receptor binding peptides, conjugated

20 compositions may be produced using recombinant DNA techniques. Briefly, a recombinant DNA molecule can be constructed which encodes both the ST receptor ligand and the toxin on a chimeric gene. When the chimeric gene is expressed, a fusion protein is produced which includes an ST receptor binding moiety and an

25 active moiety. Protein toxins are also useful to form conjugated compounds with ST receptor binding peptides through non-peptidyl bonds.

In addition, there are other approaches for utilizing active agents for the treatment of cancer. For example,

30 conjugated compositions may be produced which include an ST binding moiety and an active moiety which is an active enzyme. The ST binding moiety specifically localizes the conjugated composition to the tumor cells. An inactive prodrug which can be converted by the enzyme into an active drug is administered

35 to the patient. The prodrug is only converted to an active drug by the enzyme which is localized to the tumor. An example of an enzyme/prodrug pair includes alkaline phosphatase/

09724983 112800

etoposidephosphate. In such a case, the alkaline phosphatase is conjugated to an ST receptor binding ligand. The conjugated compound is administered and localizes at the metastasized cell. Upon contact with etoposidephosphate (the prodrug), the
5 etoposidephosphate is converted to etoposide, a chemotherapeutic drug which is taken up by the cancer cell.

Radiosensitizing agents are substances that increase the sensitivity of cells to radiation. Examples of radiosensitizing agents include nitroimidazoles, metronidazole
10 and misonidazole (see: DeVita, V.T. Jr. in *Harrison's Principles of Internal Medicine*, p.68, McGraw-Hill Book Co., N.Y. 1983, which is incorporated herein by reference). The conjugated compound that comprises a radiosensitizing agent as the active moiety is administered and localizes at the
15 metastasized cell. Upon exposure of the individual to radiation, the radiosensitizing agent is "excited" and causes the death of the cell.

Radionuclides may be used in pharmaceutical compositions that are useful for radiotherapy or imaging
20 procedures.

Examples of radionuclides useful as toxins in radiation therapy include: ^{47}Sc , ^{67}Cu , ^{90}Y , ^{109}Pd , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{199}Au , ^{211}At , ^{212}Pb and ^{212}Bi . Other radionuclides which have been used by those having ordinary skill in the art
25 include: ^{32}P and ^{33}P , ^{71}Ge , ^{77}As , ^{103}Pb , ^{105}Rh , ^{111}Ag , ^{119}Sb , ^{121}Sn , ^{131}Cs , ^{143}Pr , ^{161}Tb , ^{177}Lu , ^{191}Os , ^{193}Pt , ^{197}Hg , all beta negative and/or auger emitters. Some preferred radionuclides include: ^{90}Y , ^{131}I , ^{211}At and $^{212}\text{Pb}/^{212}\text{Bi}$.

According to the present invention, the active
30 moieties may be an imaging agent. Imaging agents are useful diagnostic procedures as well as the procedures used to identify the location of metastasized cells. Imaging can be performed by many procedures well-known to those having ordinary skill in the art and the appropriate imaging agent
35 useful in such procedures may be conjugated to an ST receptor ligand by well-known means. Imaging can be performed, for example, by radioscintigraphy, nuclear magnetic resonance

009724983 112800

imaging (MRI) or computed tomography (CT scan). The most commonly employed radionuclide imaging agents include radioactive iodine and indium. Imaging by CT scan may employ a heavy metal such as iron chelates. MRI scanning may employ
5 chelates of gadolinium or manganese. Additionally, positron emission tomography (PET) may be possible using positron emitters of oxygen, nitrogen, iron, carbon, or gallium. Example of radionuclides useful in imaging procedures include:
10 ⁴³K, ⁵²Fe, ⁵⁷Co, ⁶⁷Cu, ⁶⁷Ga, ⁶⁸Ga, ⁷⁷Br, ⁸¹Rb/^{81m}Kr, ^{87m}Sr, ^{99m}Tc, ¹¹¹In, ^{113m}In, ¹²³I, ¹²⁵I, ¹²⁷Cs, ¹²⁹Cs, ¹³¹I, ¹³²I, ¹⁹⁷Hg, ²⁰³Pb and ²⁰⁶Bi.

It is preferred that the conjugated compositions be non-immunogenic or immunogenic at a very low level. Accordingly, it is preferred that the ST receptor binding moiety be a small, poorly immunogenic or non-immunogenic
15 peptide or a non-peptide. Likewise, it is preferred that the active moiety be a small, poorly-immunogenic or non-immunogenic peptide or a non-peptide. Native ST, being a small peptide, has been shown to poorly immunogenic. (See: Klipstein, F.A. et al. (1982) *Infect. Immun.* 37:550-557; Giannella, R.A. et al.
20 (1981) *Infect. Immun.* 33:186; Burgess, M.N. et al. (1978) *Infect. Immun.* 21:60; Evans, D.G. et al. (1973) *Infect. Immun.* 7:873; Gyles, C.L. (1979) *Ann. N.Y. Acad. Sci.* 16:314; and Sack, R.B. (1975) *Ann. Rev. Microbiol.* 29:333.) Similarly, fragments and amino acid substitute derivatives of native ST
25 are poorly immunogenic. Accordingly, conjugated compositions which include all or part of the native ST as an ST receptor binding moiety are generally poorly immunogenic to the extent that the native ST is poorly immunogenic.

ST receptor ligands are conjugated to active agents
30 by a variety of well-known techniques readily performed without undue experimentation by those having ordinary skill in the art. The technique used to conjugate the ST receptor ligand to the active agent is dependent upon the molecular nature of the ST receptor ligand and the active agent. After the ST receptor
35 ligand and the active agent are conjugated to form a single molecule, assays may be performed to ensure that the conjugated molecule retains the activities of the moieties. The ST

008277 "E864260

receptor binding assay described above may be performed using the conjugated compound to test whether it is capable of binding to the ST receptor. Similarly, the activity of the active moiety may be tested using various assays for each
5 respective type of active agent. Radionuclides retain there activity, i.e. their radioactivity, irrespective of conjugation. With respect to active agents which are toxins, drugs and targeting agents, standard assays to demonstrate the activity of unconjugated forms of these compounds may be used
10 to confirm that the activity has been retained.

Conjugation may be accomplished directly between the ST receptor ligand and the active agent or linking, intermediate molecular groups may be provided between the ST receptor ligand and the active agent. Crosslinkers are
15 particularly useful to facilitate conjugation by providing attachment sites for each moiety. Crosslinkers may include additional molecular groups which serve as spacers to separate the moieties from each other to prevent either from interfering with the activity of the other.

20 In some preferred embodiments, the ST receptor ligand peptide is SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 or fragments or derivatives thereof. It has been observed that conjugation to these molecules is preferably performed at the amino terminus of each respective peptide. In ST receptor
25 ligand peptides comprising D amino acid sequences in the opposite order as SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54, conjugation preferably is performed at the carboxy terminus.

One having ordinary skill in the art may conjugate an ST receptor ligand peptide to a chemotherapeutic drug using
30 well-known techniques. For example, Magerstadt, M. *Antibody Conjugates and Malignant Disease*. (1991) CRC Press, Boca Raton, USA, pp. 110-152) which is incorporated herein by reference, teaches the conjugation of various cytostatic drugs to amino acids of antibodies. Such reactions may be applied to
35 conjugate chemotherapeutic drugs to ST receptor ligands, including ST receptor binding peptides, with an appropriate linker. ST receptor ligands which have a free amino group such

009277" 8842760

In order to conjugate an ST receptor ligand peptide to a peptide-based active agent such as a toxin, the ST receptor ligand and the toxin may be produced as a single, fusion protein either by standard peptide synthesis or recombinant DNA technology, both of which can be routinely performed by those having ordinary skill in the art. Alternatively, two peptides, the ST receptor ligand peptide and the peptide-based toxin may be produced and/or isolated as separate peptides and conjugated using crosslinkers. As with conjugated compositions that contain chemotherapeutic drugs, conjugation of ST receptor binding peptides and toxins can exploit the ability to modify the single free amino group of an

ST receptor binding peptide while preserving the receptor-binding function of this molecule.

One having ordinary skill in the art may conjugate an ST receptor ligand peptide to a radionuclide using well-known techniques. For example, Magerstadt, M. (1991) *Antibody Conjugates And Malignant Disease*, CRC Press, Boca Raton, FLA,; and Barchel, S.W. and Rhodes, B.H., (1983) *Radioimaging and Radiotherapy*, Elsevier, NY, NY, each of which is incorporated herein by reference, teach the conjugation of various therapeutic and diagnostic radionuclides to amino acids of antibodies. Such reactions may be applied to conjugate radionuclides to ST receptor ligand peptides or to ST receptor ligands including ST receptor ligand peptides with an appropriate linker.

The present invention provides pharmaceutical compositions that comprise the conjugated compounds of the invention and pharmaceutically acceptable carriers or diluents. The pharmaceutical composition of the present invention may be formulated by one having ordinary skill in the art. Suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field, which is incorporated herein by reference. In carrying out methods of the present invention, conjugated compounds of the present invention can be used alone or in combination with other diagnostic, therapeutic or additional agents. Such additional agents include excipients such as coloring, stabilizing agents, osmotic agents and antibacterial agents.

The conjugated compositions of the invention can be, for example, formulated as a solution, suspension or emulsion in association with a pharmaceutically acceptable parenteral vehicle. Examples of such vehicles are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Liposomes may also be used. The vehicle may contain additives that maintain isotonicity (e.g., sodium chloride, mannitol) and chemical stability (e.g., buffers and preservatives). The formulation is sterilized by commonly used techniques. For

003211 FEB 22 1990

example, a parenteral composition suitable for administration by injection is prepared by dissolving 1.5% by weight of active ingredient in 0.9% sodium chloride solution.

The pharmaceutical compositions according to the present invention may be administered as either a single dose or in multiple doses. The pharmaceutical compositions of the present invention may be administered either as individual therapeutic agents or in combination with other therapeutic agents. The treatments of the present invention may be combined with conventional therapies, which may be administered sequentially or simultaneously.

The pharmaceutical compositions of the present invention may be administered by any means that enables the conjugated composition to reach the targeted cells. In some embodiments, routes of administration include those selected from the group consisting of intravenous, intraarterial, intraperitoneal, local administration into the blood supply of the organ in which the tumor resides or directly into the tumor itself. Intravenous administration is the preferred mode of administration. It may be accomplished with the aid of an infusion pump.

The dosage administered varies depending upon factors such as: the nature of the active moiety; the nature of the conjugated composition; pharmacodynamic characteristics; its mode and route of administration; age, health, and weight of the recipient; nature and extent of symptoms; kind of concurrent treatment; and frequency of treatment.

Because conjugated compounds are specifically targeted to cells with ST receptors, conjugated compounds which comprise chemotherapeutics or toxins are administered in doses less than those which are used when the chemotherapeutics or toxins are administered as unconjugated active agents, preferably in doses that contain up to 100 times less active agent. In some embodiments, conjugated compounds which comprise chemotherapeutics or toxins are administered in doses that contain 10-100 times less active agent as an active moiety than the dosage of chemotherapeutics or toxins administered as

09724983 "112800

unconjugated active agents. To determine the appropriate dose, the amount of compound is preferably measured in moles instead of by weight. In that way, the variable weight of different ST binding moieties does not affect the calculation. Presuming a one to one ratio of ST binding moiety to active moiety in conjugated compositions of the invention, less moles of conjugated compounds may be administered as compared to the moles of unconjugated compounds administered, preferably up to 100 times less moles.

Typically, chemotherapeutic conjugates are administered intravenously in multiple divided doses.

Up to 20 gm IV/dose of methotrexate is typically administered in an unconjugated form. When methotrexate is administered as the active moiety in a conjugated compound of the invention, there is a 10-to 100-fold dose reduction. Thus, presuming each conjugated compound includes one molecule of methotrexate conjugated to one ST receptor binding moiety, of the total amount of conjugated compound administered, up to about 0.2 - 2.0 g of methotrexate is present and therefore administered. In some embodiments, of the total amount of conjugated compound administered, up to about 200 mg - 2g of methotrexate is present and therefore administered.

Methotrexate has a molecular weight of 455. One mole of the ST peptide-methotrexate conjugate weighs between about 1755-2955 depending on the ST peptide used. The effective dose range for ST peptide-methotrexate conjugate is between about 10 to 1000 mg. In some embodiments, dosages of 50 to 500 mg of ST peptide-methotrexate conjugate are administered. In some embodiments, dosages of 80 to 240 mg of ST peptide-methotrexate conjugate are administered.

Doxorubicin and daunorubicin each weigh about 535. Thus, ST peptide-doxorubicin conjugates and ST peptide-daunorubicin conjugates each have molecular weights of between about 1835-2553.5. Presuming each conjugated compound includes one molecule of doxorubicin or daunorubicin conjugated to one ST receptor binding moiety, the effective dose range for ST peptide-doxorubicin conjugate or ST peptide-daunorubicin

09724983-112800

conjugate is between about 40 to 4000 mg. In some embodiments, dosages of 100 to 1000 mg of ST peptide-doxorubicin conjugate or ST peptide-daunorubicin conjugate are administered. In some
5 conjugate or ST peptide-daunorubicin conjugate are administered.

Toxin-containing conjugated compounds are formulated for intravenous administration. Using this approach, up to 6 nanomoles/kg of body weight of toxin have been administered as
10 a single dose with marked therapeutic effects in patients with melanoma (Spitler L.E., et al. (1987) *Cancer Res.* 47:1717). In some embodiments, up to about 11 micrograms of ST peptide-toxin conjugated compound/kg of body weight may be administered for therapy.

15 Presuming each conjugated compound includes one molecule of ricin toxin A chain conjugated to an ST receptor binding moiety, conjugated compositions comprising ricin toxin A chain are administered in doses in which the proportion by weight of ricin toxin A chain is 1-500 μ g of the total weight
20 of the conjugated compound administered. In some preferred embodiments, conjugated compositions comprising ricin toxin A chain are administered in doses in which the proportion by weight of ricin toxin A chain is 10-100 μ g of the total weight of the conjugated compound administered. In some preferred
25 embodiments, conjugated compositions comprising ricin toxin A chain are administered in doses in which the proportion by weight of ricin toxin A chain is 2-50 μ g of the total weight of the conjugated compound administered. The molecular weight of ricin toxin A chain is 32,000. Thus, a conjugated compound
30 that contains ricin A chain linked to an ST peptide has a molecular weight of about 33,300-34,500. The range of doses of such conjugated compounds to be administered are 1 to 500 μ g. In some embodiments, 10 to 100 μ g of such conjugated compounds are administered. In some embodiments, 20 to 50 μ g of such
35 conjugated compounds are administered.

Presuming each conjugated compound includes one molecule of diphtheria toxin A chain conjugated to an ST

0924983-112800

receptor binding moiety, conjugated compositions comprising diphtheria toxin A chain are administered in doses in which the proportion by weight of diphtheria toxin A chain is 1-500 μg of the total weight of the conjugated compound administered. In some preferred embodiments, conjugated compositions comprising diphtheria toxin A chain are administered in doses in which the proportion by weight of diphtheria toxin A chain is 10-100 μg of the total weight of the conjugated compound administered. In some preferred embodiments, conjugated compositions comprising diphtheria toxin A chain are administered in doses in which the proportion by weight of diphtheria toxin A chain is 40-80 μg of the total weight of the conjugated compound administered. The molecular weight of diphtheria toxin A chain is 66,600. Thus, a conjugated compound that contains diphtheria A chain linked to an ST peptide has a molecular weight of about 67,900-69,100. The range of doses of such conjugated compounds to be administered tested are 1 to 500 μg . In some embodiments, 10 to 100 μg of such conjugated compounds are administered. In some embodiments, 40 to 80 μg of such conjugated compounds are administered.

Presuming each conjugated compound includes one molecule of *Pseudomonas* exotoxin conjugated to an ST receptor binding moiety, conjugated compositions comprising *Pseudomonas* exotoxin are administered in doses in which the proportion by weight of *Pseudomonas* exotoxin is .01 - 100 μg of the total weight of the conjugated compound administered. In some preferred embodiments, conjugated compositions comprising *Pseudomonas* exotoxin are administered in doses in which the proportion by weight of *Pseudomonas* exotoxin is .1 - 10 μg of the total weight of the conjugated compound administered. In some preferred embodiments, conjugated compositions comprising *Pseudomonas* exotoxin are administered in doses in which the proportion by weight of *Pseudomonas* exotoxin is .3 - 2.2 μg of the total weight of the conjugated compound administered. The molecular weight of *Pseudomonas* exotoxin is 22,000. Thus, a conjugated compound that contains *Pseudomonas* exotoxin linked to an ST peptide has a molecular weight of about 23,300-24,500.

05724983 112800

The range of doses of such conjugated compounds to be administered tested are .01 to 100 μ g. In some embodiments, .1 to 10 μ g of such conjugated compounds are administered. In some embodiments, .3 to 2.2 μ g of such conjugated compounds are administered.

To dose conjugated compositions comprising ST receptor binding moieties linked to active moieties that are radioisotopes in pharmaceutical compositions useful as imaging agents, it is presumed that each ST receptor binding moiety is linked to one radioactive active moiety. The amount of radioisotope to be administered is dependent upon the radioisotope. Those having ordinary skill in the art can readily formulate the amount of conjugated compound to be administered based upon the specific activity and energy of a given radionuclide used as an active moiety. Typically 0.1-100 millicuries per dose of imaging agent, preferably 1-10 millicuries, most often 2-5 millicuries are administered. Thus, pharmaceutical compositions according to the present invention useful as imaging agents which comprise conjugated compositions comprising an ST receptor binding moiety and a radioactive moiety comprise 0.1-100 millicuries, in some embodiments preferably 1-10 millicuries, in some embodiments preferably 2-5 millicuries, in some embodiments more preferably 1-5 millicuries. Examples of dosages include: ^{131}I = between about 0.1-100 millicuries per dose, in some embodiments preferably 1-10 millicuries, in some embodiments 2-5 millicuries, and in some embodiments about 4 millicuries; ^{111}In = between about 0.1-100 millicuries per dose, in some embodiments preferably 1-10 millicuries, in some embodiments 1-5 millicuries, and in some embodiments about 2 millicuries; $^{99\text{m}}\text{Tc}$ = between about 0.1-100 millicuries per dose, in some embodiments preferably 5-75 millicuries, in some embodiments 10-50 millicuries, and in some embodiments about 27 millicuries. Depending upon the specific activity of the radioactive moiety and the weight of the ST receptor binding moiety the dosage defined by weight varies. ST peptides have molecular weights of between about 1300-2500. In the

pharmaceutical composition comprising an ST peptide linked to a single ^{131}I in which the specific activity of ^{131}I -ST peptide is about 2000 Ci/mmol, administering the dose of 0.1-100 millicuries is the equivalent of 0.1-100 μg ^{131}I -ST peptide, 5 administering the dose of 1-10 millicuries is the equivalent of 1-10 μg of ^{131}I -ST peptide, administering the dose of 2-5 millicuries is equivalent to giving 2-5 μg of ^{131}I -ST peptide and administering the dose of 1-5 millicuries is equivalent to giving 1-5 μg of ^{131}I -ST peptide. In the pharmaceutical 10 composition comprising an ST peptide linked to a single ^{111}In in which the specific activity of ^{111}In -ST peptide is about 1 Ci/mmol, administering the dose of 0.1-100 millicuries is the equivalent of 0.2-200 mg ^{111}In -ST peptide, administering the dose of 1-10 millicuries is the equivalent of 2-20 mg of ^{111}In - 15 ST peptide, administering the dose of 2-5 millicuries is equivalent to giving 4-10 mg of ^{111}In -ST peptide and administering the dose of 1-5 millicuries is equivalent to giving 2-10 mg of ^{111}In -ST peptide.

To dose conjugated compositions comprising ST receptor 20 binding moieties linked to active moieties that are radioisotopes in pharmaceutical compositions useful as therapeutic agents, it is presumed that each ST receptor binding moiety is linked to one radioactive active moiety. The amount of radioisotope to be administered is dependent upon the 25 radioisotope. Those having ordinary skill in the art can readily formulate the amount of conjugated compound to be administered based upon the specific activity and energy of a given radionuclide used as an active moiety. For therapeutics that comprise ^{131}I , between 10-1000 nM, preferably 50-500, more 30 preferably about 300 nanomoles of ^{131}I at the tumor, per gram of tumor, is desirable. Thus, if there is about 1 gram of tumor, and about 0.1% of the administered dose binds to the tumor, 0.5-100 mg of ^{131}I -ST peptide conjugated compound is administered. In some embodiments, 1 to 50 mg of ^{131}I -ST 35 peptide conjugated compound is administered. In some embodiments, 5 to 10 mg of ^{131}I -ST peptide conjugated compound is administered. Wessels B.W. and R.D. Rogus (1984) *Med. Phys.*

00924983 112800

11:638 and Kwok, C.S. et al. (1985) *Med. Phys.* 12:405, both of which are incorporated herein by reference, disclose detailed dose calculations for diagnostic and therapeutic conjugates which may be used in the preparation of pharmaceutical compositions of the present invention which include radioactive conjugated compounds.

One aspect of the present invention relates to a method of treating individuals suspected of suffering from metastasized colorectal cancer. Such individuals may be treated by administering to the individual a pharmaceutical composition that comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radiostable therapeutic agent. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radiostable active agent and the ST receptor binding moiety is a peptide. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radiostable active agent and the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radiostable active agent and the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:54. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor

003211260

Ins
C2

binding moiety and an active moiety wherein the active moiety is a radiostable therapeutic agent. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radiostable active agent selected from the group consisting of: methotrexate, doxorubicin, daunorubicin, cytosinarabioside, etoposide, 5-4 fluorouracil, melphalan, chlorambucil, cis-platinum, vindesine, mitomycin, bleomycin, purothionin, macromomycin, 1,4-benzoquinone derivatives, trenimon, ricin, ricin A chain, *Pseudomonas* exotoxin, diphtheria toxin, *Clostridium perfringens* phospholipase C, bovine pancreatic ribonuclease, pokeweed antiviral protein, abrin, abrin A chain, cobra venom factor, gelonin, saporin, modeccin, viscumin, volkensin, alkaline phosphatase, nitroimidazole, metronidazole and misonidazole. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof and the active moiety is a radiostable active agent selected from the group consisting of: methotrexate, doxorubicin, daunorubicin, cytosinarabioside, etoposide, 5-4 fluorouracil, melphalan, chlorambucil, cis-platinum, vindesine, mitomycin, bleomycin, purothionin, macromomycin, 1,4-benzoquinone derivatives, trenimon, ricin, ricin A chain, *Pseudomonas* exotoxin, diphtheria toxin, *Clostridium perfringens* phospholipase C, bovine pancreatic ribonuclease, pokeweed antiviral protein, abrin, abrin A chain, cobra venom factor, gelonin, saporin, modeccin, viscumin, volkensin, alkaline phosphatase, nitroimidazole, metronidazole and misonidazole. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor

00321" E3642/60
Ins
C2

binding moiety and an active moiety wherein the active moiety is a radiostable active agent selected from the group consisting of: methotrexate, doxorubicin, daunorubicin, cytosinarabioside, cis-platin, vindesine, mitomycin and
5 bleomycin, alkaline phosphatase, ricin A chain, *Pseudomonas* exotoxin and diphtheria toxin. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an
10 active moiety wherein the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:54 and the active moiety is a radiostable active agent selected from the group consisting of: methotrexate, doxorubicin, daunorubicin,
15 cytosinarabioside, cis-platin, vindesine, mitomycin and bleomycin, alkaline phosphatase, ricin A chain, *Pseudomonas* exotoxin and diphtheria toxin. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a
20 radiostable conjugated compound described in Example 1. The individual being treated may be diagnosed as having metastasized colorectal cancer or may be diagnosed as having localized colorectal cancer and may undergo the treatment proactively in the event that there is some metastasis as yet
25 undetected. The pharmaceutical composition contains a therapeutically effective amount of the conjugated composition. A therapeutically effective amount is an amount which is effective to cause a cytotoxic or cytostatic effect on metastasized colorectal cancer cells without causing lethal
30 side effects on the individual.

One aspect of the present invention relates to a method of treating individuals suspected of suffering from metastasized colorectal cancer. Such individuals may be treated by administering to the individual a pharmaceutical
35 composition that comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the

INS
C2
09724983-112800

active moiety is a radioactive. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an

5 active moiety wherein the active moiety is a radioactive and the ST receptor binding moiety is a peptide. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor

10 binding moiety and an active moiety wherein the active moiety is a radioactive and the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof. In some

15 embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive and the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID

20 NO:5, SEQ ID NO:6 and SEQ ID NO:54. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive agent

25 selected from the group consisting of: ^{47}Sc , ^{67}Cu , ^{90}Y , ^{109}Pd , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{199}Au , ^{211}At , ^{212}Pb , ^{212}B , ^{32}P and ^{33}P , ^{71}Ge , ^{77}As , ^{103}Pb , ^{105}Rh , ^{111}Ag , ^{119}Sb , ^{121}Sn , ^{131}Cs , ^{143}Pr , ^{161}Tb , ^{177}Lu , ^{191}Os , ^{193}Mpt and ^{197}Hg . In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically

30 acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof and the active moiety is

35 a radioactive agent selected from the group consisting of: ^{47}Sc , ^{67}Cu , ^{90}Y , ^{109}Pd , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{199}Au , ^{211}At , ^{212}Pb , ^{212}B , ^{32}P and ^{33}P , ^{71}Ge , ^{77}As , ^{103}Pb , ^{105}Rh , ^{111}Ag , ^{119}Sb , ^{121}Sn , ^{131}Cs , ^{143}Pr ,

09724983-112800

161Tb, 177Lu, 191Os, 193Mpt, 197Hg, 32P and 33P, 71Ge, 77As, 103Pb, 105Rh, 111Ag, 119Sb, 121Sn, 131Cs, 143Pr, 161Tb, 177Lu, 191Os, 193Mpt, 197Hg, all beta negative and/or auger emitters. In some embodiments of the present invention, the pharmaceutical composition comprises

5 a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive agent selected from the group consisting of: 47Sc, 67Cu, 90Y, 109Pd, 123I, 125I, 131I, 186Re, 188Re, 199Au, 211At, 212Pb, 212B,

10 32P and 33P, 71Ge, 77As, 103Pb, 105Rh, 111Ag, 119Sb, 121Sn, 131Cs, 143Pr, 161Tb, 177Lu, 191Os, 193Mpt and 197Hg. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated

15 active moiety wherein the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:54 and the active moiety is a radioactive agent selected from the group consisting of: 47Sc, 67Cu, 90Y, 109Pd, 123I, 125I, 131I, 186Re, 188Re,

20 199Au, 211At, 212Pb, 212B, 32P and 33P, 71Ge, 77As, 103Pb, 105Rh, 111Ag, 119Sb, 121Sn, 131Cs, 143Pr, 161Tb, 177Lu, 191Os, 193Mpt and 197Hg. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a radioactive conjugated compound described in

25 Example 1. The individual being treated may be diagnosed as having metastasized colorectal cancer or may be diagnosed as having localized colorectal cancer and may undergo the treatment proactively in the event that there is some metastasis as yet undetected. The pharmaceutical composition

30 contains a therapeutically effective amount of the conjugated composition. A therapeutically effective amount is an amount which is effective to cause a cytotoxic or cytostatic effect on metastasized colorectal cancer cells without causing lethal side effects on the individual.

35 One aspect of the present invention relates to a method of detecting metastasized colorectal cancer cells in an individual suspected of suffering from metastasized colorectal

09724983 112800

cancer by radioimaging. Such individuals may be diagnosed as suffering from metastasized colorectal cancer and the metastasized colorectal cancer cells may be detected by administering to the individual, preferably by intravenous administration, a pharmaceutical composition that comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive and detecting the presence of a localized accumulation or aggregation of radioactivity, indicating the presence of cells with ST receptors. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive and the ST receptor binding moiety is a peptide. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive and the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive and the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:54. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive agent selected from the group consisting of: radioactive heavy metals such as iron chelates, radioactive chelates of gadolinium or manganese, positron emitters of oxygen, nitrogen, iron, carbon,

or gallium, ⁴³K, ⁵²Fe, ⁵⁷Co, ⁶⁷Cu, ⁶⁷Ga, ⁶⁸Ga, ⁷⁷Br, ⁸¹Rb/⁸¹MKr, ⁸⁷Msr, ⁹⁹Mtc, ¹¹¹In, ¹¹³MIn, ¹²³I, ¹²⁵I, ¹²⁷Cs, ¹²⁹Cs, ¹³¹I, ¹³²I, ¹⁹⁷Hg, ²⁰³Pb and ²⁰⁶Bi. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof and the active moiety is a radioactive agent selected from the group consisting of: radioactive heavy metals such as iron chelates, radioactive chelates of gadolinium or manganese, positron emitters of oxygen, nitrogen, iron, carbon, or gallium, ⁴³K, ⁵²Fe, ⁵⁷Co, ⁶⁷Cu, ⁶⁷Ga, ⁶⁸Ga, ⁷⁷Br, ⁸¹Rb/⁸¹MKr, ⁸⁷Msr, ⁹⁹Mtc, ¹¹¹In, ¹¹³MIn, ¹²³I, ¹²⁵I, ¹²⁷Cs, ¹²⁹Cs, ¹³¹I, ¹³²I, ¹⁹⁷Hg, ²⁰³Pb and ²⁰⁶Bi. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the active moiety is a radioactive agent selected from the group consisting of: ⁴³K, ⁵²Fe, ⁵⁷Co, ⁶⁷Cu, ⁶⁷Ga, ⁶⁸Ga, ⁷⁷Br, ⁸¹Rb/⁸¹MKr, ⁸⁷Msr, ⁹⁹Mtc, ¹¹¹In, ¹¹³MIn, ¹²³I, ¹²⁵I, ¹²⁷Cs, ¹²⁹Cs, ¹³¹I, ¹³²I, ¹⁹⁷Hg, ²⁰³Pb and ²⁰⁶Bi. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a conjugated compound that comprises an ST receptor binding moiety and an active moiety wherein the ST receptor binding moiety is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:54 and the active moiety is a radioactive agent selected from the group consisting of: ⁴³K, ⁵²Fe, ⁵⁷Co, ⁶⁷Cu, ⁶⁷Ga, ⁶⁸Ga, ⁷⁷Br, ⁸¹Rb/⁸¹MKr, ⁸⁷Msr, ⁹⁹Mtc, ¹¹¹In, ¹¹³MIn, ¹²³I, ¹²⁵I, ¹²⁷Cs, ¹²⁹Cs, ¹³¹I, ¹³²I, ¹⁹⁷Hg, ²⁰³Pb and ²⁰⁶Bi. In some embodiments of the present invention, the pharmaceutical composition comprises a pharmaceutically acceptable carrier or diluent and a radioactive conjugated compound described in Example 1. The individual being treated may be diagnosed as having metastasized colorectal cancer or may be diagnosed as having

localized colorectal cancer and may undergo the treatment proactively in the event that there is some metastasis as yet undetected. The pharmaceutical composition contains a diagnostically effective amount of the conjugated composition.

5 A diagnostically effective amount is an amount which can be detected at a site in the body where cells with ST receptors are located without causing lethal side effects on the individual.

Another aspect of the invention relates to unconjugated compositions which comprise an ST receptor binding ligand and an active agent. For example, liposomes are small vesicles composed of lipids. Drugs can be introduced into the center of these vesicles. The outer shell of these vesicles comprise an ST receptor binding ligand. *Liposomes* Volumes 1, 2 and 3 CRC Press Inc. Boca Raton FLA, which is incorporated herein by reference, disclose preparation of liposome-encapsulated active agents which include targeting agents that correspond to ST receptor ligand in the outer shell. Unconjugated compositions which comprise an ST receptor ligand in the matrix of a liposome with an active agent inside include such compositions in which the ST receptor ligand is selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof and the active agent is selected from the group consisting of: methotrexate, doxorubicin, daunorubicin, cytosinarabioside, etoposide, 5-fluorouracil, melphalan, chlorambucil, cis-platinum, vindesine, mitomycin, bleomycin, purothionin, macromomycin, 1,4-benzoquinone derivatives, trenimon, ricin, ricin A chain, *Pseudomonas* exotoxin, diphtheria toxin, *Clostridium perfringens* phospholipase C, bovine pancreatic ribonuclease, pokeweed antiviral protein, abrin, abrin A chain, cobra venom factor, gelonin, saporin, modeccin, viscumin, volkensin, alkaline phosphatase, nitroimidazole, metronidazole and misonidazole.

Another aspect of the invention relates to
35 unconjugated and conjugated compositions which comprise an ST
receptor ligand used to deliver therapeutic nucleic acid
molecules to cells that comprise an ST receptor such as normal

5

20

35

presence in the cell inhibits production of an undesirable protein. In some embodiments, the nucleic acid molecule encodes a ribozyme whose presence in the cell inhibits production of an undesirable protein. In some embodiments, the nucleic acid molecule encodes a protein or peptide that is desirably produced in the cell. In some embodiments, the nucleic acid molecule encodes a functional copy of a gene that is defective in the targeted cell. The nucleic acid molecule is preferably operably linked to regulatory elements needed to express the coding sequence in the cell.

Liposomes are small vesicles composed of lipids. Genetic constructs which encode proteins that are desired to be expressed in ST receptor-bearing cells are introduced into the center of these vesicles. The outer shell of these vesicles
15 comprise an ST receptor ligand, in some embodiments preferably an ST peptide. *Liposomes* Volumes 1, 2 and 3 CRC Press Inc. Boca Raton FLA, which is incorporated herein by reference, disclose preparation of liposome-encapsulated active agents which include antibodies in the outer shell. In the present
20 invention, an ST receptor ligand such as for example an ST peptide corresponds to the antibodies in the outer shell. Unconjugated compositions which comprise an ST receptor ligand in the matrix of a liposome with an active agent inside include such compositions in which the ST receptor ligand is selected
25 from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 and fragments and derivatives thereof.

In one embodiment for example, cystic fibrosis, a genetic disease in which there is a mutation of a specific gene encoding a chloride transport protein which ultimately produces abnormalities of function in many systems, most notably in the respiratory and intestinal tract, is treated by gene therapy techniques using ST receptor ligands to deliver the corrective gene to cells. Current therapy has been directed at replacing the mutant gene in the respiratory system with the normal gene by targeting these genes directly to the cells lining the respiratory tract using viruses which bind only to those cells. Similarly, the normal gene is packaged in liposomes targeted on

their surface with ST receptor ligands and delivered to the intestinal tract. ST receptor ligands specifically target and direct the liposomes containing the normal gene to correct the lesion for cystic fibrosis to the specific cells lining the intestinal tract, from the duodenum to the rectum. Uptake of that genetic material by those cells should result in a cure of cystic fibrosis in the intestinal tract.

In another embodiment, the delivery of normal copies of the p53 tumor suppressor gene to the intestinal tract is accomplished using ST receptor ligand to target the gene therapeutic. Mutations of the p53 tumor suppressor gene appears to play a prominent role in the development of colorectal cancer in the intestinal tract. One approach to combatting this disease is the delivery of normal copies of this gene to the intestinal tract to cells expressing mutant forms of this gene. Genetic constructs that comprise normal p53 tumor suppressor genes are incorporated into liposomes that comprise an ST receptor ligand. The composition is delivered to the intestinal tract. ST receptor binding ligands specifically target and direct the liposomes containing the normal gene to correct the lesion created by mutation of p53 suppressor gene in intestinal cells.

Preparation of genetic constructs is with the skill of those having ordinary skill in the art. The present invention allows such construct to be specifically targeted by using the ST receptor ligands of the present invention. The compositions of the invention include an ST receptor ligand such as an ST peptide associated with a delivery vehicle and a gene construct which comprises a coding sequence for a protein whose production is desired in the cells of the intestinal tract linked to necessary regulatory sequences for expression in the cells. For uptake by cells of the intestinal tract, the compositions are administered orally or by enema whereby they enter the intestinal tract and contact cells which comprise ST receptors. The delivery vehicles associate with the ST receptor by virtue of the ST receptor ligand and the vehicle is internalized into the cell or the active agent/genetic

construct is otherwise taken up by the cell. Once internalized, the construct can provide a therapeutic effect on the individual. One having ordinary skill in the art can readily formulate such compositions for oral or enema
5 administration and determine the effective amount of such composition to be administered to treat the disease or disorder.

The following examples are illustrative but are not meant to be limiting of the present invention.

10 Examples

Example 1

The following are representative compounds according to the present invention. Whenever stated below, reference to a series of compounds is provided for efficiency and is meant
15 to name each compound in the series including all the compounds in numerical order, such as for example "3-D1 to 3-D16" is meant to refer to compounds 3-D1, 3-D2, 3-D3, 3-D4, 3-D5, 3-D6, 3-D7, 3-D8, 3-D9, 3-D10, 3-D11, 3-D12, 3-D13, 3-D14, 3-D15 and 3-D16. Likewise, whenever stated below, reference to a series
20 of SEQ ID NO:'s is provided for efficiency and is meant to name each SEQ ID NO: in the series including the all SEQ ID NO:'s in numerical order, such as for example SEQ ID NO:5 through SEQ ID NO:54 is meant to refer to SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ
25 ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID
30 NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53 and SEQ ID NO:54. Similarly, whenever
35 stated below, reference to a series of compounds is provided for efficiency and is meant to name each compound in the series

including the all compounds in numerical order, such as for example "5-AP to 54-AP" is meant to refer to compounds 5-AP, 6-AP, 7-AP, 8-AP, 9-AP, 10-AP, 11-AP, 12-AP, 13-AP, 14-AP, 15-AP, 16-AP, 17-AP, 18-AP, 19-AP, 20-AP, 21-AP, 22-AP, 23-AP, 24-AP, 5 25-AP, 26-AP, 27-AP, 28-AP, 29-AP, 30-AP, 31-AP, 32-AP, 33-AP, 34-AP, 35-AP, 36-AP, 37-AP, 38-AP, 39-AP, 40-AP, 41-AP, 42-AP, 43-AP, 44-AP, 45-AP, 46-AP, 47-AP, 48-AP, 49-AP, 50-AP, 51-AP, 52-AP, 53-AP and 54-AP.

Compound 2-D1 comprises methotrexate (amethopterin)
10 conjugated to SEQ ID NO:2.

Compound 2-D2 comprises doxorubicin (adrimycin)
conjugated to SEQ ID NO:2.

Compound 2-D3 comprises daunorubicin conjugated to SEQ
ID NO:2.

15 Compound 2-D4 comprises cytosinarabinoside conjugated
to SEQ ID NO:2.

Compound 2-D5 comprises etoposide conjugated to SEQ
ID NO:2.

20 ~~Compound 2-D6 comprises 5-4 fluorouracil conjugated~~
to SEQ ID NO:2.

Compound 2-D7 comprises melphalan conjugated to SEQ
ID NO:2.

Compound 2-D8 comprises chlorambucil conjugated to SEQ
ID NO:2.

25 Compound 2-D9 comprises cyclophosphamide conjugated
to SEQ ID NO:2.

Compound 2-D10 comprises cis-platinum conjugated to
SEQ ID NO:2.

30 Compound 2-D11 comprises vindesine conjugated to SEQ
ID NO:2.

Compound 2-D12 comprises mitomycin conjugated to SEQ
ID NO:2.

Compound 2-D13 comprises bleomycin conjugated to SEQ
ID NO:2.

35 Compound 2-D14 comprises purothionin conjugated to SEQ
ID NO:2.

0082112800

Ins
C4

Compound 2-D16 comprises trenimon conjugated to SEQ
ID NO:2.

Compounds 5-D1 to 5-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 5-D1 to 5-D16 each comprise SEQ ID NO:5 as the ST receptor binding moiety.

Compounds 7-D1 to 7-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 7-D1 to 7-D16 each comprise SEQ ID NO:7 as the ST receptor binding moiety.

25 Compounds 9-D1 to 9-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 9-D1 to 9-D16 each comprise SEQ ID NO:9 as the ST receptor binding moiety.

Compounds 12-D1 to 12-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 12-D1 to

12-D16 each comprise SEQ ID NO:11 as the ST receptor binding moiety.

Compounds 12-D1 to 12-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ
5 ID NO:2 as the ST receptor binding moiety, compounds 12-D1 to 12-D16 each comprise SEQ ID NO:12 as the ST receptor binding moiety.

Compounds 13-D1 to 13-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ
10 ID NO:2 as the ST receptor binding moiety, compounds 13-D1 to 13-D16 each comprise SEQ ID NO:13 as the ST receptor binding moiety.

Compounds 14-D1 to 14-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ
15 ID NO:2 as the ST receptor binding moiety, compounds 14-D1 to 14-D16 each comprise SEQ ID NO:14 as the ST receptor binding moiety.

Compounds 15-D1 to 15-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ
20 ID NO:2 as the ST receptor binding moiety, compounds 15-D1 to 15-D16 each comprise SEQ ID NO:15 as the ST receptor binding moiety.

Compounds 16-D1 to 16-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ
25 ID NO:2 as the ST receptor binding moiety, compounds 16-D1 to 16-D16 each comprise SEQ ID NO:16 as the ST receptor binding moiety.

Compounds 17-D1 to 17-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ
30 ID NO:2 as the ST receptor binding moiety, compounds 17-D1 to 17-D16 each comprise SEQ ID NO:17 as the ST receptor binding moiety.

Compounds 18-D1 to 18-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ
35 ID NO:2 as the ST receptor binding moiety, compounds 18-D1 to 18-D16 each comprise SEQ ID NO:18 as the ST receptor binding moiety.

003211 084260

Compounds 19-D1 to 19-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 19-D1 to 19-D16 each comprise SEQ ID NO:19 as the ST receptor binding moiety.

Compounds 20-D1 to 20-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 20-D1 to 20-D16 each comprise SEQ ID NO:20 as the ST receptor binding moiety.

Compounds 22-D1 to 22-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 22-D1 to 22-D16 each comprise SEQ ID NO:21 as the ST receptor binding moiety.

Compounds 22-D1 to 22-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 22-D1 to 22-D16 each comprise SEQ ID NO:22 as the ST receptor binding moiety.

Compounds 23-D1 to 23-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 23-D1 to 23-D16 each comprise SEQ ID NO:23 as the ST receptor binding moiety.

Compounds 24-D1 to 24-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 24-D1 to 24-D16 each comprise SEQ ID NO:24 as the ST receptor binding moiety.

Compounds 25-D1 to 25-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 25-D1 to 25-D16 each comprise SEQ ID NO:25 as the ST receptor binding moiety.

Compounds 26-D1 to 26-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ

00827 112800

ID NO:2 as the ST receptor binding moiety, compounds 26-D1 to 26-D16 each comprise SEQ ID NO:26 as the ST receptor binding moiety.

Compounds 27-D1 to 27-D16 are the same as compounds
5 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 27-D1 to 27-D16 each comprise SEQ ID NO:27 as the ST receptor binding moiety.

Compounds 28-D1 to 28-D16 are the same as compounds
10 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 28-D1 to 28-D16 each comprise SEQ ID NO:28 as the ST receptor binding moiety.

Compounds 29-D1 to 29-D16 are the same as compounds
15 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 29-D1 to 29-D16 each comprise SEQ ID NO:29 as the ST receptor binding moiety.

Compounds 30-D1 to 30-D16 are the same as compounds
20 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 30-D1 to 30-D16 each comprise SEQ ID NO:30 as the ST receptor binding moiety.

Compounds 32-D1 to 32-D16 are the same as compounds
25 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 32-D1 to 32-D16 each comprise SEQ ID NO:31 as the ST receptor binding moiety.

Compounds 32-D1 to 32-D16 are the same as compounds
30 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 32-D1 to 32-D16 each comprise SEQ ID NO:32 as the ST receptor binding moiety.

Compounds 33-D1 to 33-D16 are the same as compounds
35 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 33-D1 to

00321" 2364260

33-D16 each comprise SEQ ID NO:33 as the ST receptor binding moiety.

Compounds 34-D1 to 34-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 34-D1 to 34-D16 each comprise SEQ ID NO:34 as the ST receptor binding moiety.

Compounds 35-D1 to 35-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 35-D1 to 35-D16 each comprise SEQ ID NO:35 as the ST receptor binding moiety.

Compounds 36-D1 to 36-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 36-D1 to 36-D16 each comprise SEQ ID NO:36 as the ST receptor binding moiety.

Compounds 37-D1 to 37-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 37-D1 to 37-D16 each comprise SEQ ID NO:37 as the ST receptor binding moiety.

Compounds 38-D1 to 38-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 38-D1 to 38-D16 each comprise SEQ ID NO:38 as the ST receptor binding moiety.

Compounds 39-D1 to 39-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 39-D1 to 39-D16 each comprise SEQ ID NO:39 as the ST receptor binding moiety.

Compounds 40-D1 to 40-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 40-D1 to 40-D16 each comprise SEQ ID NO:40 as the ST receptor binding moiety.

009724983 112800

5

10

15

20

25

30

35

Compounds 48-D1 to 48-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ

ID NO:2 as the ST receptor binding moiety, compounds 48-D1 to 48-D16 each comprise SEQ ID NO:48 as the ST receptor binding moiety.

5 Compounds 49-D1 to 49-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 49-D1 to 49-D16 each comprise SEQ ID NO:49 as the ST receptor binding moiety.

10 Compounds 50-D1 to 50-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 50-D1 to 50-D16 each comprise SEQ ID NO:50 as the ST receptor binding moiety.

15 Compounds 51-D1 to 51-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 51-D1 to 51-D16 each comprise SEQ ID NO:51 as the ST receptor binding moiety.

20 Compounds 52-D1 to 52-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 52-D1 to 52-D16 each comprise SEQ ID NO:52 as the ST receptor binding moiety.

25 Compounds 53-D1 to 53-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 53-D1 to 53-D16 each comprise SEQ ID NO:53 as the ST receptor binding moiety.

30 Compounds 54-D1 to 54-D16 are the same as compounds 2-D1 to 2-D16, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 54-D1 to 54-D16 each comprise SEQ ID NO:54 as the ST receptor binding moiety.

35 Compound 2-T1 comprises ricin conjugated to SEQ ID NO:2.

Compound 2-T2 comprises ricin A chain (ricin toxin) conjugated to SEQ ID NO:2.

00827T 0354260

Compound 2-T3 comprises *Pseudomonas* exotoxin (PE) conjugated to SEQ ID NO:2.

Compound 2-T4 comprises diphtheria toxin (DT), conjugated to SEQ ID NO:2.

5 Compound 2-T5 comprises *Clostridium perfringens* phospholipase C (PLC) conjugated to SEQ ID NO:2.

Compound 2-T6 comprises bovine pancreatic ribonuclease (BPR) conjugated to SEQ ID NO:2.

10 Compound 2-T7 comprises pokeweed antiviral protein (PAP) conjugated to SEQ ID NO:2.

Compound 2-T8 comprises abrin conjugated to SEQ ID NO:2.

Compound 2-T9 comprises abrin A chain (abin toxin) conjugated to SEQ ID NO:2.

15 Compound 2-T10 comprises cobra venom factor (CVF) conjugated to SEQ ID NO:2.

Compound 2-T11 comprises gelonin (GEL) conjugated to SEQ ID NO:2.

20 Compound 2-T12 comprises saporin (SAP) conjugated to SEQ ID NO:2.

Compound 2-T13 comprises modeccin conjugated to SEQ ID NO:2.

Compound 2-T14 comprises viscumin conjugated to SEQ ID NO:2.

25 Compound 2-T15 comprises volkensin conjugated to SEQ ID NO:2.

30 Compounds 3-T1 to 3-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 3-T1 to 3-T15 each comprise SEQ ID NO:3 as the ST receptor binding moiety.

Compounds 5-T1 to 5-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 5-T1 to 5-T15 each comprise SEQ ID NO:5 as the ST receptor binding moiety.

35 Compounds 6-T1 to 6-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID

00827T" E864260

NO:2 as the ST receptor binding moiety, compounds 6-T1 to 6-T15 each comprise SEQ ID NO:6 as the ST receptor binding moiety.

Compounds 7-T1 to 7-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 7-T1 to 7-T15 each comprise SEQ ID NO:7 as the ST receptor binding moiety.

Compounds 8-T1 to 8-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 8-T1 to 8-T15 each comprise SEQ ID NO:8 as the ST receptor binding moiety.

Compounds 9-T1 to 9-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 9-T1 to 9-T15 each comprise SEQ ID NO:9 as the ST receptor binding moiety.

Compounds 10-T1 to 10-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 10-T1 to 10-T15 each comprise SEQ ID NO:2 as the ST receptor binding moiety.

Compounds 11-T1 to 11-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 11-T1 to 11-T15 each comprise SEQ ID NO:11 as the ST receptor binding moiety.

Compounds 12-T1 to 12-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 12-T1 to 12-T15 each comprise SEQ ID NO:12 as the ST receptor binding moiety.

Compounds 13-T1 to 13-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 13-T1 to 13-T15 each comprise SEQ ID NO:13 as the ST receptor binding moiety.

Compounds 14-T1 to 14-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 14-T1 to

008277" 0864260

14-T15 each comprise SEQ ID NO:14 as the ST receptor binding moiety.

Compounds 15-T1 to 15-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 15-T1 to 15-T15 each comprise SEQ ID NO:15 as the ST receptor binding moiety.

Compounds 15-T1 to 15-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 15-T1 to 15-T15 each comprise SEQ ID NO:15 as the ST receptor binding moiety.

Compounds 17-T1 to 17-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 17-T1 to 17-T15 each comprise SEQ ID NO:17 as the ST receptor binding moiety.

Compounds 18-T1 to 18-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 18-T1 to 18-T15 each comprise SEQ ID NO:18 as the ST receptor binding moiety.

Compounds 19-T1 to 19-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 19-T1 to 19-T15 each comprise SEQ ID NO:19 as the ST receptor binding moiety.

Compounds 20-T1 to 20-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 20-T1 to 20-T15 each comprise SEQ ID NO:20 as the ST receptor binding moiety.

Compounds 21-T1 to 21-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 21-T1 to 21-T15 each comprise SEQ ID NO:21 as the ST receptor binding moiety.

003277" E8642460

5

10

15

20

25

30

35

2

ID NO:2 as the ST receptor binding moiety, compounds 29-T1 to 29-T15 each comprise SEQ ID NO:29 as the ST receptor binding moiety.

Compounds 30-T1 to 30-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 30-T1 to 30-T15 each comprise SEQ ID NO:30 as the ST receptor binding moiety.

Compounds 31-T1 to 31-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 31-T1 to 31-T15 each comprise SEQ ID NO:31 as the ST receptor binding moiety.

Compounds 32-T1 to 32-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 32-T1 to 32-T15 each comprise SEQ ID NO:32 as the ST receptor binding moiety.

Compounds 33-T1 to 33-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 33-T1 to 33-T15 each comprise SEQ ID NO:33 as the ST receptor binding moiety.

Compounds 34-T1 to 34-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 34-T1 to 34-T15 each comprise SEQ ID NO:34 as the ST receptor binding moiety.

Compounds 35-T1 to 35-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 35-T1 to 35-T15 each comprise SEQ ID NO:35 as the ST receptor binding moiety.

Compounds 36-T1 to 36-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 36-T1 to

00827" 08642460

36-T15 each comprise SEQ ID NO:36 as the ST receptor binding moiety.

Compounds 37-T1 to 37-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 37-T1 to 37-T15 each comprise SEQ ID NO:37 as the ST receptor binding moiety.

Compounds 38-T1 to 38-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 38-T1 to 38-T15 each comprise SEQ ID NO:38 as the ST receptor binding moiety.

Compounds 39-T1 to 39-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 39-T1 to 39-T15 each comprise SEQ ID NO:39 as the ST receptor binding moiety.

Compounds 40-T1 to 40-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 40-T1 to 40-T15 each comprise SEQ ID NO:40 as the ST receptor binding moiety.

Compounds 41-T1 to 41-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 41-T1 to 41-T15 each comprise SEQ ID NO:41 as the ST receptor binding moiety.

Compounds 42-T1 to 42-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 42-T1 to 42-T15 each comprise SEQ ID NO:42 as the ST receptor binding moiety.

Compounds 43-T1 to 43-T15 are the same as compounds 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 43-T1 to 43-T15 each comprise SEQ ID NO:43 as the ST receptor binding moiety.

SECRET

5

10

15

20

25

30

35

2

ID NO:2 as the ST receptor binding moiety, compounds 51-T1 to 51-T15 each comprise SEQ ID NO:51 as the ST receptor binding moiety.

Compounds 52-T1 to 52-T15 are the same as compounds 5 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 52-T1 to 52-T15 each comprise SEQ ID NO:52 as the ST receptor binding moiety.

Compounds 53-T1 to 53-T15 are the same as compounds 10 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 53-T1 to 53-T15 each comprise SEQ ID NO:53 as the ST receptor binding moiety.

Compounds 54-T1 to 54-T15 are the same as compounds 15 2-T1 to 2-T15, respectively, except instead of comprising SEQ ID NO:2 as the ST receptor binding moiety, compounds 54-T1 to 54-T15 each comprise SEQ ID NO:54 as the ST receptor binding moiety.

Compounds 2-AP, 3-AP and 5-AP to 54-AP refer to the 20 51 conjugated compounds that comprise alkaline phosphatase conjugated to SEQ ID NO:2, SEQ ID NO:3 and SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-NIZ, 3-NIZ and 5-NIZ to 54-NIZ refer to the 51 conjugated compounds that comprise nitroimidazole 25 conjugated to SEQ ID NO:2, SEQ ID NO:3 and SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-MEZ, 3-MEZ and 5-MEZ to 54-MEZ refer to the 51 conjugated compounds that comprise metronidazole conjugated to SEQ ID NO:2, SEQ ID NO:3 and SEQ ID NO:5 through 30 SEQ ID NO:54, respectively.

Compounds 2-MIS, 3-MIS and 5-MIS to 54-MIS refer to the 51 conjugated compounds that comprise misonidazole conjugated to SEQ ID NO:2, SEQ ID NO:3 and SEQ ID NO:5 through 35 SEQ ID NO:54, respectively.

Compounds 2-47Sc, 3-47Sc and 5-47Sc to 54-47Sc refer to the 51 conjugated compounds that comprise ⁴⁷Sc conjugated to

00821" 0964260

SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-67Cu, 3-67Cu and 5-67Cu to 54-67Cu refer to the 51 conjugated compounds that comprise ⁶⁷Cu conjugated to
5 SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-90Y, 3-90Y and 5-90Y to 54-90Y refer to the 51 conjugated compounds that comprise ⁹⁰Y conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54,
10 respectively.

Compounds 2-109Pd, 3-109Pd and 5-109Pd to 54-109Pd refer to the 51 conjugated compounds that comprise ¹⁰⁹Pd conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-123I, 3-123I and 5-123I to 54-123I refer to the 51 conjugated compounds that comprise ¹²³I conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-125I, 3-125I and 5-125I to 54-125I refer to the 51 conjugated compounds that comprise ¹²⁵I conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-131I, 3-131I and 5-131I to 54-131I refer to the 51 conjugated compounds that comprise ¹³¹I conjugated to
25 SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-132I, 3-132I and 5-132I to 54-132I refer to the 51 conjugated compounds that comprise ¹³²I conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54,
30 respectively.

Compounds 2-186Re, 3-186Re and 5-186Re to 54-186Re refer to the 51 conjugated compounds that comprise ¹⁸⁶Re, conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-188Re, 3-188Re and 5-188Re to 54-188Re refer to the 51 conjugated compounds that comprise ¹⁸⁸Re,

00321" 0854260

conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through
SEQ ID NO:54, respectively.

Compounds 2-199Au, 3-199Au and 5-199Au to 54-199Au refer to the 51 conjugated compounds that comprise ¹⁹⁹Au, 5 conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-211At, 3-211At and 5-211At to 54-211At refer to the 51 conjugated compounds that comprise ²¹¹At, conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through
10 SEQ ID NO:54, respectively.

Compounds 2-212Pb, 3-212Pb and 5-212Pb to 54-212Pb refer to the 51 conjugated compounds that comprise ²¹²Pb conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

15 Compounds 2-212Bi, 3-212Bi and 5-212Bi to 54-212Bi
refer to the 51 conjugated compounds that comprise ²¹²Bi
conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through
SEQ ID NO:54, respectively.

Compounds 2-203Pb, 3-203Pb and 5-203Pb to 54-203Pb
20 refer to the 51 conjugated compounds that comprise ²⁰³Pb
conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through
SEQ ID NO:54, respectively.

Compounds 2-206Bi, 3-206Bi and 5-206Bi to 54-206Bi refer to the 51 conjugated compounds that comprise ²⁰⁶Bi conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-32P, 3-32P and 5-32P to 54-32P refer to the 51 conjugated compounds that comprise ³²P conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-33P, 3-33P and 5-33P to 54-33P refer to the 51 conjugated compounds that comprise ³³P conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

35 Compounds 2-⁷¹Ge, 3-⁷¹Ge and 5-⁷¹Ge to 54-⁷¹Ge refer
to the 51 conjugated compounds that comprise ⁷¹Ge conjugated to

SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-77As, 3-77As and 5-77As to 54-77As refer to the 51 conjugated compounds that comprise ⁷⁷As conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-103Pd, 3-103Pd and 5-103Pd to 54-103Pd refer to the 51 conjugated compounds that comprise ¹⁰³Pd conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through 10 SEQ ID NO:54, respectively.

Compounds 2-105Rh, 3-105Rh and 5-105Rh to 54-105Rh refer to the 51 conjugated compounds that comprise ¹⁰⁵Rh conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

15 Compounds 2-111Ag, 3-111Ag and 5-111Ag to 54-111Ag
refer to the 51 conjugated compounds that comprise ¹¹¹Ag
conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through
SEQ ID NO:54, respectively.

Compounds 2-119Sb, 3-119Sb and 5-119Sb to 54-119Sb
20 refer to the 51 conjugated compounds that comprise ¹¹⁹Sb
conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through
SEQ ID NO:54, respectively.

Compounds 2-121Sn, 3-121-Sn and 5-121Sn to 54-121Sn refer to the 51 conjugated compounds that comprise ¹²¹Sn
25 conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through
SEQ ID NO:54, respectively.

Compounds 2-131Cs, 3-131Cs and 5-131Cs to 54-131Cs refer to the 51 conjugated compounds that comprise ¹³¹Cs conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through 30 SEQ ID NO:54, respectively.

Compounds 2-127Cs, 3-131Cs and 5-131Cs to 54-127Cs refer to the 51 conjugated compounds that comprise ¹²⁷Cs conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

35 Compounds 2-129Cs, 3-129Cs and 5-129Cs to 54-129Cs
refer to the 51 conjugated compounds that comprise ¹²⁹Cs

0272493160

Compounds 2-143Pr, 3-143Pr and 5-143Pr to 54-143Pr refer to the 51 conjugated compounds that comprise ¹⁴³Pr conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-177Lu, 3-177Lu and 5-177Lu to 54-177Lu refer to the 51 conjugated compounds that comprise ¹⁷⁷Lu conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-193mPt, 3-193mPt and 5-193mPt to 54-193mPt refer to the 51 conjugated compounds that comprise ¹⁹³mPt conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-43K, 3-43K and 5-43K to 54-43K refer to the 51 conjugated compounds that comprise ⁴³K conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

35 Compounds 2-⁵⁷Co, 3-⁵⁷Co and 5-⁵⁷Co to 54-⁵⁷Co refer
to the 51 conjugated compounds that comprise ⁵⁷Co conjugated to

SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-67Ga, 3-67Ga and 5-67Ga to 54-67Ga refer to the 51 conjugated compounds that comprise ⁶⁷Ga conjugated to
5 SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-68Ga, 3-68Ga and 5-68Ga to 54-68Ga refer to the 51 conjugated compounds that comprise ⁶⁸Ga conjugated to
10 SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-77Br, 3-77Br and 5-77Br to 54-77Br refer to the 51 conjugated compounds that comprise ⁷⁷Br conjugated to
SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-81Rb, 3-81Rb and 5-81Rb to 54-81Rb refer to the 51 conjugated compounds that comprise ⁸¹Rb conjugated to
15 SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-81mKr, 3-81mKr and 5-81mKr to 54-81mKr
20 refer to the 51 conjugated compounds that comprise ^{81m}Kr conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-87mSr, 3-87mSr and 5-87mSr to 54-87mSr refer to the 51 conjugated compounds that comprise ^{87m}Sr
25 conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-99mTc, 3-99mTc and 5-99mTc to 54-99mTc refer to the 51 conjugated compounds that comprise ^{99m}Tc
30 conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-111In, 3-111In and 5-111In to 54-111In refer to the 51 conjugated compounds that comprise ¹¹¹In
conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

Compounds 2-113mIn, 3-113mIn and 5-113mIn to 54-113mIn refer to the 51 conjugated compounds that comprise ^{113m}In
35

003211 11200 092493 11200

conjugated to SEQ ID NO:2, SEQ ID NO:3 AND SEQ ID NO:5 through SEQ ID NO:54, respectively.

The compounds described in this example are combined with a pharmaceutically acceptable carrier or diluent to produce pharmaceutical compositions according to the present invention. Radiostable compounds described herein are useful in pharmaceutical compositions as therapeutics in the treatment of individuals suspected of suffering from metastasized colorectal cancer including treatment of individuals diagnosed with localized colorectal cancer as a prophylactic/therapeutic before metastasis can be readily detected. When present in therapeutically effective amounts, radioactive compounds described herein are useful in pharmaceutical compositions as therapeutic agents in the treatment of individuals suspected of suffering from metastasized colorectal cancer including treatment of individuals diagnosed with localized colorectal cancer as a prophylactic/therapeutic before metastasis can be readily detected. When present in diagnostically effective amounts, radioactive compounds described herein are useful in pharmaceutical compositions as imaging agents in the diagnosis and identification of metastasized colorectal cancer in individuals.

Example 2

One procedure for crosslinking ST receptor ligands which have a free amino group such as ST receptor binding peptides, as for example SEQ ID NO:2, SEQ ID NO:3, and SEQ ID NOS:5-54 to active agents which have a free amino group such as methotrexate, doxorubicin, daunorubicin, cytosinarabioside, cis-platin, vindesine, mitomycin and bleomycin, or alkaline phosphatase, or protein- or peptide-based toxin employs homobifunctional succinimidyl esters, preferably with chain carbon spacers such as disuccinimidyl suberate (Pierce Co, Rockford, IL). This approach of amino group derivatization has been employed successfully to crosslink native ST to biotin and, ultimately, to large agarose beads of micron-scale size, preserving the function of native ST (Hughes, M., et al. (1991)

Biochem. 30:10738; Hakki, S., et al. (1993) *Int. J. Biochem.* 25:557; Almenoff, J.S., et al. (1992) *Mol. Micro.* 8:865; each of which is incorporated herein by reference).

An ST binding ligand with the free amino group such as an ST receptor binding peptide is incubated in the presence of the chemical crosslinking agent and an active agent which have a free amino group in equimolar quantities at room temperature for 15-30 min. Incubation is terminated by separating the reactants by gel permeation chromatography by HPLC. This technique separates the conjugated compounds from free active agents and free ST binding ligands, active agent-active agent conjugates and ST binding ligand-ST binding ligand conjugates. Homogeneous preparations of conjugated through their free amino groups and with a preferred molar ratio of 1:1 are obtained. As indicated above, complexing the free amino group of an ST peptide preserves receptor binding function.

Example 3

In the event that a cleavable conjugated compound is required, the same protocol as described above may be employed utilizing 3,3'- dithiobis (sulfosuccinimidylpropionate (SPDP); Pierce, IL.). SPDP forms a sulfhydryl group from a free amino group which may be used to conjugate a compound to another free amino group. For example, ST peptides such as SEQ ID NO:2, SEQ ID NO:3, SEQ ID NOS:5-54 are derivatized using established procedures employing N-succinimidyl-3 (2-pyridyldithio)-propionate (SPDP, Pharmacia-LKB, NJ). The ST peptide is incubated with a 5-fold molar excess of SPDP for 30 minutes at room temperature. The ST-pyridylthiopropionate conjugate is separated from unreacted reagents by gel permeation chromatography by HPLC. An active agent with a free amino group, such as a protein-based toxin, is prepared for conjugation by reduction with dithiothreitol for 4 hours at room temperature. Reduced active agent is incubated with a 2-fold molar excess of ST receptor ligand-PDP conjugate at pH 8.0 for 36 hours at 4°C. Conjugate compound is purified from unreacted agents by gel permeation chromatography by HPLC.

This protocol for conjugation is particularly useful to conjugate ST peptides to diphtheria toxin A chains and *Pseudomonas* exotoxin as well as ricin toxin A chains (Magerstadt, M. *Antibody Conjugates and Malignant Disease*. (1991) CRC Press, Boca Raton, USA, pp. 110-152; Cawley, D.B. et al. (1980) *Cell* 22:563; Cumber, A.J., et al. (1985) *Meth. Enz.* 112:207; Gros, O. (1985) *J. Immunol. Meth.* 81:283; Worrell, N.R., et al. (1986) *Anti-Cancer Drug Design* 1:179; Thorpe, P.E. et al. (1987) *Cancer Res.* 47:5924, each of which is incorporated herein by reference).

Example 4

Active agents with a free amino group may be derivatized with SPDP as described above and conjugated with an ST ligand that has a free amino group and that has been modified with the succinimidyl ester of iodoacetic acid (Pierce Co., Rockford, IL) (Magerstadt, M. (1991) *Antibody Conjugates And Malignant Disease*, CRC Press Boca Raton; Cumber, A.J. et al. (1985) *Meth. Enz.* 112:20, which are incorporated herein by reference). Conjugation relies on the selective reaction of iodoacetyl groups introduced into the amino terminal of the ST ligand with the thiol groups introduced into the active agent. As with the above protocol, this procedure avoids homopolymer formation. However, the product is conjugated through a central thioether linkage which cannot be reduced.

Example 5

An ST receptor ligand with a free amino group and active agents with free amino groups may be conjugated through a disulfide bond using iminothiolane (Pierce, Rockford, IL) (Fitzgerald, D.J.P. et al. (1983) *Cell* 32:607; Magerstadt, M. (1991) *Antibody Conjugates And Malignant Disease*, CRC Press, Boca Raton; Bjorn, M.J., et al. (1985) *Cancer Res.* 45:1214; Bjorn, M.J., et al. (1986) *Cancer Res.* 46:3262, which are incorporated herein by reference). The ST receptor ligand with a free amino group is derivatized at the amino terminal with iminothiolane and the active agent is derivatized with SPDP as

described above. Reacting iminothiolane-derivatized ST receptor ligand with SPDP-derivatized active agent results in conjugation by a reducible disulfide bond. In addition, iminothiolane provides the versatility to conjugate these
5 proteins through bonds other than disulfides. Thus, derivatization of active agents with the heterobifunctional agent sulfosuccinimidyl 4-(N-maleimidomethyl) cyclohexane (Pierce, Rockford, IL) and reaction with iminothiolane-derivatized ST receptor ligand will conjugate these peptides
10 without formation of disulfides.

Example 6

Conjugated compounds according to the invention which comprise an active moiety that is a therapeutic agent specifically inhibit T84 cells *in vitro*. The following
15 protocols may be used to demonstrate that the conjugated compounds according to the invention which comprise an active moiety that is chemotherapeutic or toxin specifically inhibit T84 cells *in vitro*. Inhibition of T84 cells is assessed by determining the effects of conjugated compounds on the ability
20 of T84 cells to incorporate ³⁵S-leucine into protein, ³H-thymidine into DNA, and to form colonies. The assessment of protein and DNA synthesis are classical techniques to determine the cytotoxicity of conjugated compounds *in vitro*. Inhibition of protein synthesis is measured because the toxins used as
25 active moieties are specific inhibitors of this process. Therefore, these assays are the most sensitive measure of whether conjugated compounds are binding to and internalized into T84 cells. Inhibition of DNA synthesis is measured because some chemotherapeutics inhibit DNA synthesis and
30 further, it is a cytotoxicity assay which correlates closely with the reproductive survivability of cells in culture. Cytotoxicity, or the disruption of normal cellular metabolic processes, may not always directly correlate with cell survivability. Therefore, assessment of colony formation will
35 directly measure the ability of the experimental agents to decrease the survivability of tumor cells, which closely correlates with the impact of therapeutic agents on tumor

00321T 2864260

5 compound. The results obtained in the test assays and control assays are compared.

10 monolayer culture by established protocols (Wilson, A.P. (1987)
"Cytotoxicity and viability assays", Animal Cell Culture: A
Practical Approach. Freshney, R.I., ed. pp. 183-216, IRL
Press, Oxford. which is incorporated herein by reference).

15 inhibit protein synthesis *in vitro*, cells are plated in 200 μ l
of medium at a sub-confluent density of $1-2 \times 10^5$ and allowed
to attach to form a dividing cell monolayer over 12 hours at
37°C. Subsequently, the media is replaced with 200 μ l of fresh
media containing the appropriate concentration of conjugated
20 compounds and cells incubated at 37°C for various amounts of
time. At the end of the indicated incubation period, cells is
washed twice with medium and incubated at 37°C in 0.5 ml of
methionine-free medium supplemented with 0.5 μ Ci of L³⁵S-
methionine (800 Ci/mmol). After incubation for another 2 hours
25 at 37°C, the medium is aspirated, cells washed twice with
medium containing 1 mg/ml of methionine, and then precipitated
in 12% ice-cold TCA. Radioactivity recovered in TCA
precipitates by centrifugation is quantified by liquid
scintillation spectroscopy. In these studies, cells are
30 maintained in log growth and assays are performed using
triplicate wells. Data is expressed as a percentage of protein
synthesis observed in the presence of experimental agents
compared to untreated cells.

35 inhibit DNA synthesis in vitro cells are plated as a subconfluent monolayer and incubated with experimental agents as described above. At the end of the incubation period, cells

are washed twice and incubated at 37°C in medium containing 2.5 μ Ci of 3 H-thymidine (5 Ci/mmol). After incubation for another hour, cells are processed with TCA, precipitates recovered, and radioactivity quantified as described above. As above, cells
5 are maintained in log growth and assays is performed in triplicate. Data is expressed as a percentage of DNA synthesis observed in the presence of experimental agents compared to untreated cells.

To assess the ability of a conjugated compound to
10 inhibit survival and proliferation by measuring colony formation in monolayer culture, cells are plated as a sub-confluent monolayer on 25 cm² flasks and allowed to attach as described above. The medium is replaced with that containing various concentrations of experimental agents and incubated
15 with cells for various amounts of time. At the end of the incubation, cells are recovered as a single cell suspension by trypsinization and replated to a density which will yield 100-200 colonies per 6 cm plate. Cells are permitted to grow for 7 days, then fixed in methanol, stained with 1% crystal violet,
20 and the number of colonies quantified. Assays are performed in duplicate and data is expressed as a percentage of colony formation observed in the presence of experimental agents compared to untreated cells. Results in our laboratory have demonstrated that T84 cells can be placed into single cell
25 suspensions utilizing trypsin (10 μ g/ml) with a plating efficiency of 40% and a doubling time of 18 hours.

Example 7

Radioactive iodine such as 123 I, 125 I, 131 I and 132 I, can be added to an ST receptor binding peptide such as an ST
30 peptide using a standard protocol well-known to those having ordinary skill in the art (Thompson, M. et al. (1985) *Analytical Biochemistry* 148:26, which is incorporated herein by reference). Radioactive iodine is conjugated directly to an ST peptide such as SEQ ID NO:2, SEQ ID NO:3 or SEQ ID NO:5 at
35 tyrosine-5, tyrosine-4 or tyrosine-5, respectively.

Briefly, the ST peptide is produced in bacteria. For example, *E. coli* strain 431 is grown in culture and secretes ST

008217" E8642460

5 which is incorporated herein by reference.

10 min after which the products are subjected to chromatography on a Sepak reversed-phase cartridge (Millipore Corp., MA) followed by separation on a C₁₈ reversed-phase column by HPLC using a 20-25% acetonitrile gradient. Conjugated compositions which comprise SEQ ID NO:2, SEQ ID NO:3 or SEQ ID NO:5 with the 15 radioiodine attached to tyrosine-4 elutes at 45 min. These molecules retain full biochemical and pharmacological activity.

Example 8

¹²⁵I is conjugated directly to an ST peptide such as
SEO ID NO:13 at tyrosine-4.

20 SEQ ID NO:13 is produced by solid-phase synthesis as described above. Ten micrograms of SEQ ID NO:13 are reacted with 2 milliCuries of ^{125}I Na (Amersham Corporation, Massachusetts) in the presence of Iodobeads (Bio Rad Laboratories, CA) and beta-D-glucose. These are reacted for 30
25 min after which the products are subjected to chromatography on a Sepak reversed-phase cartridge (Millipore Corp., MA) followed by separation on a C_{18} reversed-phase column by HPLC using a 20-25% acetonitrile gradient. ^{125}I -SEQ ID NO:13 conjugate with the radioiodine attached to tyrosine-4 elutes at 45 min. This
30 molecule retains full biochemical and pharmacological activity.

Dosing of radioiodine for diagnostic imaging typically requires about 4 milliCuries/patient (Steinstraber, A., et al. (1988) *J. Nucl. Med.* 29:875; Wessels, B.W. and Rogus, R.D. (1984) *Med. Phys.* 11:638; Kwok, C.S., et al. (1985) *Med. Phys.* 12:405). For proteins labeled with a specific activity of 2,000 Curies/mmol, such as ST peptide, this would require about

Example 9

SEQ ID NO:13 is produced by solid-phase synthesis as described above. Ten micrograms of SEQ ID NO:13 are reacted with 10 milliCuries of ¹³¹INa (Amersham Corporation, Massachusetts) in the presence of Iodobeads (Bio Rad Laboratories, CA) and beta-D-glucose. These are reacted for 30 min after which the products are subjected to chromatography on a Sepak reversed-phase cartridge (Millipore Corp., MA) followed by separation on a C₁₈ reversed-phase column by HPLC using a 20-25% acetonitrile gradient. ¹³¹I-SEQ ID NO:13 conjugate with the radioiodine attached to tyrosine-4 elutes at 45 min. This molecule retains full biochemical and pharmacological activity.

25 **Example 10**

In some embodiments, coupling of ST receptor ligands which have a free amino group, particularly ST receptor binding peptides such as ST peptides, and active agents with a free amino group such as protein-based toxins is performed by introducing a disulfide bridge between the 2 molecules. This strategy is particularly useful to conjugate ST peptides since the free amino terminal has been shown to be useful as a point of conjugation without affecting ST binding activity. This strategy is particularly useful to conjugate protein-based toxins since the free amino terminal is available on such molecules and for some conjugated compounds, most notably RTA conjugates, a disulfide bridge which can be reduced to yield

al. (1983) *Infec. Immun.* 42:539, which is incorporated herein by reference).

Toxins are prepared for coupling by reduction with 0.1 M dithiothreitol (DTT) for 4 hours at room temperature in 0.4 M Tris-HCl, pH 8.0 and 1mM EDTA. Reduced toxins are desalted on a Sephadex G-25 column equilibrated in TES buffer and mixed with a 2-fold molar excess of ST-PDP. Reactions are adjusted to pH 8.0 with TES and incubated at 4°C for 36 hours. ST peptide-toxin conjugates are purified from unreacted products and homopolymers of ST peptides and toxins by gel filtration on Sephadex G-75 in 20 mM TES, pH 8.0 containing 0.1 M NaCl. Chromatographic fractions are monitored by SDS-PAGE on 10% polyacrylamide gels under non-reducing conditions for the presence of 1:1 conjugates of ST peptides and toxins. Also, these conjugates are analyzed by 10% SDS-PAGE under reducing conditions, to insure that ST and cytotoxins are coupled by a reducible disulfide bond. Molar concentrations of the conjugate are calculated by quantifying radioactivity in these samples.

ST trace labeled with ^{125}I on tyrosine 4 (10 Ci/mmol) is used in order to follow the conjugate through various separation and chromatographic steps and to enable us to calculate the molar ratio of ST to cytotoxin in the final purified conjugate. ST trace labeled with ^{125}I is derivatized by incubating 1 mg/ml with a 5-fold molar excess of SPDP for 30 min at room temperature in Na phosphate buffer, pH 7.4. The ST-pyridylthiopropionate (ST-PDP) conjugate is purified from unreacted crosslinking agent by chromatography on Sephadex G-25 equilibrated with 20 mM N-Tris (hydroxymethyl)-methyl-2-aminoethane sulfonic acid (TES) buffer, pH 7.4. Preservation of receptor binding of conjugated ST peptides in human intestinal membranes is determined in competition assays of increasing concentrations of ST-PDP and ^{125}I -ST ($5 \times 10^{10}\text{M}$), to insure that this process does not destroy the function of the ST receptor ligand.

The above coupling protocol has several advantages for conjugating the various toxins. First, it introduces a

00821" 112800

reducible disulfide bridge into the conjugated composition, important for RTA cytotoxicity. Also, this technique avoids the exposure of ST peptide to quantitative reduction with DTT which could interrupt its 3 intrachain disulfide bonds
5 important for receptor binding activity. In addition, there is a single group available at the amino terminal of ST peptide for derivatization with SPDP and previous experiments have demonstrated that derivatization of that group preserves the binding properties of the ligand. Therefore, other
10 configurations for conjugation which could result in inactivation of ST are not possible. Furthermore, PEA requires preactivation with DTT to achieve optimum cytotoxicity which will be accomplished utilizing the above protocol.

To produce a functional conjugated compound that
15 comprises a toxin, it is essential that the receptor binding and enzyme activities of the moieties are preserved throughout the process of conjugation. Therefore, once such conjugate compounds are obtained, they are tested for the preservation of those functions. ST receptor binding activity of conjugated
20 compounds is examined in competitive binding assays, as described above. In these studies, increasing concentrations of the conjugated compounds are incubated with a constant concentration ($5 \times 10^{10}M$) of ^{125}I -ST and intestinal membranes (50-100 μg of protein) to achieve equilibrium. Parallel
25 incubations contain excess (5×10^7M) unlabeled ST to assess non-specific binding. The concentration-dependent competitive displacement of radiolabeled ST by conjugated compounds is compared to the competitive displacement achieved by native ST. Displacement curves are employed to estimate the affinity of
30 each conjugated compound (K_D) and compare that to the affinity of native ST measured by this technique. Control studies include evaluating the ability of unconjugated toxins to compete with native ST for receptor binding. These studies establish that the binding function of ST in the conjugated
35 construct is preserved.

Preservation of toxin activity in conjugated compounds is also assessed. PEA and DTA induce toxicity by catalyzing

008277" E8642460

the NAD-dependent ADP-ribosylation of elongation factor 2 (EF2), inhibiting protein synthesis. ADP-ribosyl transferase activity is assessed using an established assay (Chung, D.W. and Collier, R.J. *Infect. Immun.* 16:832; Fitzgerald, D.J.P. (1987) *Meth. Enz.* 151:139, which are both incorporated herein by reference). Reactions are conducted in 30 Mm Tris-HCl, pH 8.2 containing 40 mM DTT, 50 mCi ¹⁴C-NAD, and 20 μl of rabbit reticulocyte lysate containing elongation factor 2 (EF-2; Promega, Madison, WI) in a total volume of 500 μl. Reactions are initiated by the addition of lysate, incubated for 30 minutes at 37°C, and terminated by the addition of ice-cold 12% TCA. Radioactivity in protein precipitates collected by centrifugation is quantified by liquid scintillation spectroscopy. The ability of the conjugated compounds that comprise DTA or PEA to catalyze the transfer of labeled ADP-ribose to EF-2 is compared to that catalyzed by similar quantities of unconjugated toxins. Control experiments include examining the ability of unconjugated toxins or ST to catalyze ADP-ribose transfer and the effects of ST on the enzymatic activity of unconjugated cytotoxins.

RTA inhibits protein synthesis by catalytically inactivating the 60S ribosomal subunit. The catalytic activity of conjugated compounds that comprise RTA is assessed by its ability to inhibit protein synthesis in cell-free assays using established procedures (Leonard, J.E. et al. (1985) *Cancer Res.* 45:5263 which is incorporated herein by reference). Assays contain 35 μl of nuclease-treated rabbit reticulocyte lysates, 1 μl of 1mM mixed amino acids deficient in methionine, 2 μl of Brome mosaic RNA (Promega, Madison, WI) at 0.5 μg/μl, 7 μl of sterile water or conjugate solution, and 5 μCi of ³⁵S-methionine in a total volume of 50 μl. Reactions will be initiated by the addition of lysate, incubated at 30°C for 30 minutes, and terminated by the use of addition of 12% TCA. Radioactivity in protein precipitates collected by centrifugation is quantified by liquid scintillation spectroscopy. Control experiments include examining the ability of unconjugated RTA or ST peptide to inhibit cell-free protein synthesis and the effects of ST

09724983-112800

peptide on the inhibitory activity of the unconjugated cytotoxin.

Example 11

Methotrexate is linked to SEQ ID NO:12 by the
5 homobifunctional crosslinker succinimidyl esters with long
chain carbon spacers such as disuccinimidyl suberate (Pierce,
IL). SEQ ID NO:12 is incubated in the presence of the chemical
crosslinking agent and methotrexate in equimolar quantities at
room temperature for 15-30 min. Incubation is terminated by
10 separating the reactants by gel permeation chromatography by
HPLC. This technique separates the methotrexate/SEQ ID NO:12
conjugates from free drug, free ST peptide, drug-drug
conjugates and ST peptide-ST peptide conjugates. Homogeneous
preparations of SEQ ID NO:12-methotrexate conjugates coupled
15 through their free amino groups and with a preferred molar
ratio of 1:1 are obtained. Complexing the free amino group of
ST preserves receptor binding function.

Example 12

¹¹¹In is coupled to SEQ ID NO:37 with functional amino
20 groups using a chelator. The ST peptide has a free amino
function at the amino terminal which may be modified without
altering the ST receptor binding activity of the ST peptide.
¹¹¹In is rapidly and potently chelated by either EDTA
(ethylenediaminetetraacetic acid) or DTPA
25 (diethylenetriaminepentaacetic acid). DTPA is preferred over
EDTA because the latter may be more unstable *in vivo*. The
¹¹¹In-DTPA is converted to a mixed N-hydroxysuccinimide ester
which is reactive with free amino groups, mixed with ST, and
the reaction products, including ¹¹¹In-SEQ ID NO:37 separated by
30 HPLC (Bremer, K.H. and Schwarz, A. (1987) in *Safety And
Efficacy Of Radiopharmaceuticals*. Kristensen, K. and
Norbygaard, E., Eds. Martinus Nijhoff, Dordrecht, The
Netherlands, P. 43; Krejcarek, G.E., and Tucker, K.L. (1977)
Biochem. Biophys. Res. Commun. 77:581; Paxton, R.J., et al.
35 (1985) *Cancer Res.* 45:5694; Richardson, A.P., et al. (1986)
Nucl. Med. Biol. 14:569, which are each incorporated herein by
reference).

003211 12300

Example 13

^{99m}Tc can be conjugated to SEQ ID NO:46 using an approach which is similar to that for indium. Thus, technetium can be chelated by DTPA which is converted to an anhydride, such as N-hydroxysuccinimide anhydride, and reacted with SEQ ID NO:46. The ST-technetium conjugate can then be separated using HPLC (Magerstadt, M. (1991) *Antibody Conjugates And Malignant Disease* CRC Press, Boca Raton; Eckelman, W.C. and Paik, C.H. (1986) *Nucl. Med. Biol.* 14:569)

Example 14

Diphtheria toxin A chain (DTA) is prepared from native diphtheria toxin by standard techniques. SEQ ID NO:22 is coupled to N-succinimidyl-3(2-pyridyldithio)-propionate (SPDP, Pharmacia-LKB, Piscataway, NJ) and the SEQ ID NO:22-PDP conjugate is purified by HPLC by established procedures. DTA is reduced with dithiothreitol and incubated with SEQ ID NO:22-PDP. DTA-SEQ ID NO:22 is purified after conjugation using HPLC.

Example 15

Pseudomonas Exotoxin is prepared from native sources by standard techniques. SEQ ID NO:54 is coupled to N-succinimidyl-3(2-pyridyldithio)-propionate (SPDP, Pharmacia-LKB, Piscataway, NJ) and the SEQ ID NO:54-PDP conjugate is purified by HPLC by established procedures. *Pseudomonas* Exotoxin is reduced with dithiothreitol and incubated with SEQ ID NO:54-PDP. *Pseudomonas* Exotoxin-SEQ ID NO:54 is purified after conjugation using HPLC.

Example 16

Doxorubicin is linked to SEQ ID NO:54 by the homobifunctional crosslinker succinimidyl esters with long chain carbon spacers such as disuccinimidyl suberate (Pierce, IL). SEQ ID NO:54 is incubated in the presence of the chemical crosslinking agent and doxorubicin in equimolar quantities at room temperature for 15-30 min. Incubation is terminated by separating the reactants by gel permeation chromatography by HPLC. This technique separates the doxorubicin/SEQ ID NO:54 conjugates from free doxorubicin, free ST peptide, drug-drug

conjugates and ST peptide-ST peptide conjugates. Homogeneous preparations of SEQ ID NO:54-doxorubicin conjugates coupled through their free amino groups and with a preferred molar ratio of 1:1 are obtained. Complexing the free amino group of
5 ST preserves receptor binding function.

Example 17

Daunorubicin is linked to SEQ ID NO:32 by the homobifunctional crosslinker succinimidyl esters with long chain carbon spacers such as disuccinimidyl suberate (Pierce,
10 IL). SEQ ID NO:32 is incubated in the presence of the chemical crosslinking agent and daunorubicin in equimolar quantities at room temperature for 15-30 min. Incubation is terminated by separating the reactants by gel permeation chromatography by HPLC. This technique separates the daunorubicin/SEQ ID NO:54
15 conjugates from free daunorubicin, free ST peptide, drug-drug conjugates and ST peptide-ST peptide conjugates. Homogeneous preparations of SEQ ID NO:54-daunorubicin conjugates coupled through their free amino groups and with a preferred molar ratio of 1:1 are obtained. Complexing the free amino group of
20 ST preserves receptor binding function.

003211 E8642460